arXiv:0809.4446v1 [math.GT] 25 Sep 2008

BOUNDED COHOMOLOGY AND l_1-HOMOLOGY OF THREE-MANIFOLDS

PIERRE DERBEZ

ABSTRACT. In this paper we define, for each aspherical orientable 3-manifold M endowed with a *torus splitting* $\mathcal{T}$, a 2-dimensional fundamental l_1-class $[M]^{\mathcal{T}}$ whose l_1-norm has similar properties as the Gromov simplicial volume of M (additivity under torus splittings and isometry under finite covering maps). Next, we use the Gromov simplicial volume of M and the l_1-norm of $[M]^{\mathcal{T}}$ to give a complete characterization of those nonzero degree maps $f\colon\ M \to N$ which are homotopic to a $\deg(f)$-covering map. As an application we characterize those degree one maps $f\colon\ M \to N$ which are homotopic to a homeomorphism in terms of bounded cohomology classes.

1. INTRODUCTION

In order to characterize those nonzero degree maps $f\colon\ M \to N$ between closed orientable aspherical 3-manifolds which are homotopic to a finite covering we first need the simplicial volume of Gromov. Recall that this invariant is obtained as follows: consider the l_1-norm on the real singular chains which induces, taking the infimum for all cycles representing a homology class, a l_1 semi-norm on the homology groups of a manifold. Then the simplicial volume of M is the l_1 semi-norm of the fundamental class $[M]$ corresponding to the orientation of M. This invariant is necessary since for any finite covering maps $p\colon\ \widetilde{M} \to M$ then $(*)$

$$\|\widetilde{M}\| = |\deg(p)|\|M\|$$

Equivalently, this equality means that p induces an isometry $p_\sharp\colon\ H_3(\widetilde{M};\mathbf{R}) \to H_3(M;\mathbf{R})$ with respect to the l_1 semi-norm. For hyperbolic 3-manifolds equality $(*)$ is sufficient to characterize covering maps. More precisely, it follows from the Perelman geometrization of 3-manifolds and from Gromov and Thurston's works that if M is a closed orientable hyperbolic 3-manifold then any nonzero degree map $f\colon\ M \to N$ into a closed orientable irreducible 3-manifold is homotopic to a covering map if and only if $f_\sharp\colon\ H_3(M;\mathbf{R}) \to H_3(N;\mathbf{R})$ is an isometry. However this condition is not sufficient to characterize covering maps between non-hyperbolic 3-manifolds. Roughly speaking this comes from the fact that the Gromov simplicial volume does not detect the "non-hyperbolic part" of an aspherical 3-manifold. Thus one of the purposes of this paper is to construct for each aspherical oriented closed 3-manifold $(M,\mathcal{T}_M)$, where $\mathcal{T}_M$ denotes the JSJ-family of canonical tori of M, a kind of *secondary fundamental class* of M denoted by $[M]^{\mathcal{T}_M}$ which detects the non-hyperboic part of M. This class lies in the second l_1-homology group of M denoted by $H_2^{l_1}(M;\mathbf{R})$ and endowed with the l_1 semi-norm $\|.\|_1$. The l_1-norm of $[M]^{\mathcal{T}_M}$ together with the Gromov simplicial volume allow to characterize finite covering maps. More precisely, the main result of this paper states as follows:

Date: September 25, 2008.

1991 *Mathematics Subject Classification.* 57M50, 51H20.

Key words and phrases. Aspherical 3-manifolds, bounded cohomology, l_1-homology, non-zero degree maps, topological rigidity.

Theorem 1.1. *Let $f\colon M \to N$ denote a nonzero degree map from an aspherical closed oriented 3-manifold into an irreducible closed oriented 3-manifold. If M is not a $\widetilde{\mathrm{SL}}(2,\mathbf{R})$-manifold then f is homotopic to a $\deg(f)$-covering map iff $\|f_\sharp([M])\|_1 = \|[M]\|_1$ and $\|f_\sharp([M]^{\mathcal{T}_M})\|_1 = \|[M]^{\mathcal{T}_M}\|_1$ where $\mathcal{T}_M$ denotes the JSJ-splitting of M.*

As a consequence of Theorem 1.1 we deduce, using the duality between l_1-homology and bounded cohomology, a complete description of those degree one maps which are homotopic to a homeomorphism which answers to a question of M. Boileau. Denote by $H^2_b(M;\mathbf{R})$ the second bounded cohomology group of M endowed with the semi-norm $\|.\|_\infty$ and by $\widehat{H}^2_b(M;\mathbf{R})$, resp. $\widehat{H}^{l_1}_2(M;\mathbf{R})$ the quotient space $H^2_b(M;\mathbf{R})/\ker\|.\|_\infty$, resp. $H^{l_1}_2(M;\mathbf{R})/\ker\|.\|_1$.

Corollary 1.2. *A degree one map $f\colon M \to N$ between closed Haken manifolds is homotopic to a homeomorphism iff $\|M\| = \|N\|$ and f induces an isometric isomorphism $f^\sharp\colon (\widehat{H}^2_b(N;\mathbf{R}),\|.\|_\infty) \to (\widehat{H}^2_b(M;\mathbf{R}),\|.\|_\infty)$, resp. an isometry $f_\sharp\colon (\widehat{H}^{l_1}_2(M;\mathbf{R}),\|.\|_1) \to (\widehat{H}^{l_1}_2(N;\mathbf{R}),\|.\|_1)$.*

Recall that when M admits a geometry $\mathbf{H}^3$ then M. Gromov and W.P. Thurston gave a characterization of local isometry using the Gromov simplicial volume. When M admits a geometry $\widetilde{\mathrm{SL}}(2,\mathbf{R})$ then Y. Rong solved this problem using the Seifert volume in [Ro1, Corollary 5.1] and when M admits a geometry $\mathbf{R}^3$, Nil or Sol then S. Wang showed in [W] that any nonzero degree map $f\colon M \to N$ is homotopic to a covering map. On the other hand, when M is a surface bundle over $\mathbf{S}^1$ (which covers the case where M admits a $\mathbf{H}^2 \times \mathbf{R}$-geometry) then M. Boileau and S. Wang gave a characterization of finite covering maps in terms of isometries with respect to the Thurston's norm (see [BW]).

2. FILLING ISOMETRIES AND THE SECONDARY FUNDAMENTAL CLASS

In this section we define the objects and we state the intermediate results which will be used in the proof of Theorem 1.1 and Corollary 1.2.

First of all recall recall that the l_1-homology groups of a topological space X are obtained as follows. Denote by $C^{l_1}_*(X)$ the l_1-completion of the real singular chains $C_*(X)$. This means that

$$C^{l_1}_*(X) = \left\{ c = \sum_{i=1}^{\infty} a_i\sigma_i \text{ s.t. } \|c\|_1 = \sum_{i=1}^{\infty} |a_i| < \infty \right\}$$

where $a_i \in \mathbf{R}$ and $\sigma_i\colon \Delta_* \to X$ is a singular simplex. We will denote by $S_*(X)$ the set of singular simplices. The topological dual of $C^{l_1}_*(X)$ is given by the set

$$C^*_b(X) = \left\{ w \in C^*(X) \text{ s.t. } \|w\|_\infty = \sup_{\sigma\in S_*(X)} |\langle w,\sigma\rangle| < \infty \right\}$$

Note that the ∂ and δ operators are bounded so that $(C^{l_1}_*(X),\partial)$ and $(C^*_b(X),\delta)$ are chain, resp. cochain, complexes. We denote by $H^{l_1}_*(X)$, resp. by $H^*_b(X)$, the homology, resp. cohomology, of this chain, resp. cochain, complex. The vector spaces $H^{l_1}_*(X)$ and $H^*_b(X)$ are endowed with the quotient semi-norm that we denote still by $\|.\|_1$ and $\|.\|_\infty$ respectively. Since a bounded operator has not necessarily a closed image then the above semi-norms are not norms in general. Thus it will be convenient to consider the reduced l_1-homology and bounded cohomology groups defined by $\widehat{H}^{l_1}_*(X) = \ker\partial/\overline{\mathrm{Im}(\partial)} =$

$H_*^{l_1}(X)/\ker \|.\|_1$ and $\widehat{H}_b^*(X) = \ker\delta/\overline{\mathrm{Im}(\delta)} = H_b^*(X)/\ker\|.\|_\infty$. The evaluation map gives Kronecker product

$$\langle .,. \rangle\colon\ H_b^*(X) \otimes H_*^{l_1}(X) \to \mathbf{R}$$

which descends to a Kronecker product on reduced groups by the Holder inequality

$$\langle .,. \rangle\colon\ \widehat{H}_b^*(X) \otimes \widehat{H}_*^{l_1}(X) \to \mathbf{R}$$

2.1. **A filling isometry.** We begin with the construction of a filling homomorphism.

Lemma 2.1. *Let (X, Y, A) be a triple of spaces where $Y \subset X$ (may be $X = Y$) and each component of $A \subset Y$ has an amenable fundamental group.*

(i) For any relative 2-cycle z in (Y, A) there exists $u \in C_2^{l_1}(A)$ such that $z + u \in Z_2^{l_1}(Y)$.

(ii) Filling homomorphism. The map $z \mapsto z + u$ induces a filling homomorphism

$$\Theta_Y\colon\ H_2(Y, A) \to H_2^{l_1}(Y) \to H_2^{l_1}(X)$$

defined by $\Theta_Y([z]) = [z + u]$.

(iii) Filling contraction. The filling homomorphism $\Theta_Y\colon\ H_2(Y, A) \to H_2^{l_1}(X)$ satisfies $\|\Theta_Y\| \le 1$ when $H_2(Y, A)$ is endowed with the l_1 semi-norm.

(iv) Naturality. Let $f\colon\ (X, Y, A) \to (Z, W, B)$ be a continuous map of triple of spaces, where each component of A and B has amenable fundamental group. Then the following diagram is consistant.

$$\begin{tikzcd}
H_2(Y, A; \mathbf{R}) \arrow[r, "\Theta_Y"] \arrow[d, "f_\sharp"'] & H_2^{l_1}(X; \mathbf{R}) \arrow[d, "f_\sharp"'] \\
H_2(W, B; \mathbf{R}) \arrow[r, "\Theta_W"] & H_2^{l_1}(Z; \mathbf{R})
\end{tikzcd}$$

In particular this result implies that Θ_Y induces a contraction homomorphism between reduced homology groups

$$\widehat{\Theta}_Y\colon\ \widehat{H}_2(Y, A) = \frac{H_2(Y, A)}{\|.\|_1} \to \widehat{H}_2^{l_1}(X)$$

Let M denote a closed aspherical orientable 3-manifold and let $\mathcal{T}$ denote a family of incompressible tori in M. We say that $\mathcal{T}$ is a *torus splitting* of M if each component of $M \setminus \mathcal{T}$ is either a Seifert manifold or has a hyperbolic interior. From now on we assume that M is endowed with a torus splitting $\mathcal{T}$. Let $(P, \partial P)$ be a component of $M \setminus \mathcal{T}$. Let $\alpha \in H_2(P, \partial P; \mathbf{R})$. Since each component of ∂P has an amenable fundamental group then we can consider the filling homomorphism

$$\widehat{\Theta}_P\colon\ \widehat{H}_2(P, \partial P; \mathbf{R}) \to \widehat{H}_2^{l_1}(M; \mathbf{R})$$

and the subspace of $H_2^{l_1}(M; \mathbf{R})$

$$\widehat{H}_2^{l_1}\left(M^{\mathcal{T}}; \mathbf{R}\right) = \mathrm{Vect}\left\langle \mathrm{Im}(\widehat{\Theta}_P) | P \text{ is a Seifert component of } M \setminus \mathcal{T} \right\rangle$$

Suppose that P admits a Seifert fibration whose fiber is denoted by h. By a *horizontal surface* we mean a properly embedded incompressible surface F in P which is transverse to h. Note that if F is a horizontal surface in P then the Seifert bundle η induces an orbifold covering $\eta|F\colon\ F \to \mathcal{O}_P$ whose degree is denoted by $d_F \neq 0$, where $\mathcal{O}_P$ denotes the base 2-orbifold. A horizontal surface will be termed *minimal* if $|d_F|$ is minimal over all horizontal surfaces in P.

Lemma 2.2. *Let (P,h) be an aspherical oriented Seifert manifold with $\mathbf{S}^1$-fiber h endowed with a fixed orientation $\mathfrak{o}(h)$. If P is either Euclidean,* Nil *or a $\widetilde{\mathrm{SL}}(2,\mathbf{R})$-manifold then $\widehat{H}_2(P,\partial P;\mathbf{R})=\{0\}$. If the interior of P admits a $\mathbf{H}^2\times\mathbf{R}$ structure then $\widehat{H}_2(P,\partial P;\mathbf{R})\simeq\mathbf{R}$. Moreover, in the latter case, if F and F' are minimal surfaces in P then $[F]=[F']$ in $\widehat{H}_2(P,\partial P;\mathbf{R})$ (where $[F]$, resp. $[F']$, corresponds to the orientation class of F, resp. F', so that $\mathfrak{o}(F)\times\mathfrak{o}(h)$, resp. $\mathfrak{o}(F')\times\mathfrak{o}(h)$, matches the given orientation of P) and $\widehat{H}_2(P,\partial P;\mathbf{R})=\langle[F]\rangle$.*

Notation. For an oriented $\mathbf{H}^2\times\mathbf{R}$-manifold P we will denote by α_P the class of any minimal surface in $\widehat{H}_2(P,\partial P;\mathbf{R})$ with the convention for orientations of Lemma 2.2. For Euclidean, Nil or a $\widetilde{\mathrm{SL}}(2,\mathbf{R})$-manifold then we set $\alpha_P=0$. We denote by $\widehat{H}_2^+(P,\partial P;\mathbf{R})$ the set defined by

$$\{\alpha\in\widehat{H}_2(P,\partial P;\mathbf{R})|\ \alpha=\xi.\alpha_P,\ \xi\geq 0\}$$

When M denotes an oriented aspherical 3-manifold endowed with a torus splitting $\mathcal{T}$ each Seifert component P of $M\setminus\mathcal{T}$ is oriented by M and we fix an orientation $\mathfrak{o}(h)$ for the fiber h of each P. In this case we say that M is *framed*. We denote by $\widehat{H}_2^{l_1,+}(M^{\mathcal{T}};\mathbf{R})$ the set defined by

$$\{\alpha\in\widehat{H}_2^{l_1}(M^{\mathcal{T}};\mathbf{R})|\ \alpha=\sum\xi_i.\widehat{\Theta}_{P_i}(\alpha_{P_i}),\ \xi_i\geq 0\}$$

where the P_i's run over the Seifert components of $M\setminus\mathcal{T}$. Notice that since the only aspherical Seifert fibered spaces admitting at least two non-isotopic fibrations are the Euclidean manifolds then the above notions are well-defined by Lemma 2.2. Some geometric properties of the map $\widehat{\Theta}_P$ are reflected in the following

Theorem 2.3. *Let $(M,\mathcal{T})$ be a closed aspherical orientable 3-manifold endowed with a torus splitting and denote by $P_1,...,P_l$ the Seifert components of $M\setminus\mathcal{T}$. Then*

(i) Isometry: the filling homomorphism $\widehat{\Theta}_{P_i}\colon\ \widehat{H}_2(P_i,\partial P_i;\mathbf{R})\to\widehat{H}_2^{l_1}(M;\mathbf{R})$ is an isometry with respect to the l_1-norms. Moreover for any $\alpha\in\widehat{H}_2(P_i,\partial P_i;\mathbf{R})$ then

$$\|\alpha\|_1=\|\widehat{\Theta}_{P_i}(\alpha)\|_1=|\xi_\alpha|\|F_i\|$$

where ξ_α is the real number such that $\alpha=\xi_\alpha.\alpha_{P_i}$ and F_i is a minimal surface in P_i. In particular, if α is represented by an incompressible connected surface $\mathcal{F}$ then

$$\|\alpha\|_1=\|\widehat{\Theta}_{P_i}(\alpha)\|_1=\|\mathcal{F}\|$$

(ii) Additivity: for any l-uple $(\alpha_1,...,\alpha_l)\in\widehat{H}_2^+(P_1,\partial P_1;\mathbf{R})\times...\times\widehat{H}_2^+(P_l,\partial P_l;\mathbf{R})$ we have

$$\|\widehat{\Theta}_{P_1}(\alpha_1)+...+\widehat{\Theta}_{P_l}(\alpha_l)\|_1=\|\widehat{\Theta}_{P_1}(\alpha_1)\|_1+...+\|\widehat{\Theta}_{P_l}(\alpha_l)\|_1$$

We end this section with a result which describes the metric behavior of finite covering maps. To this purpose note that throughout this paper we adopt the following convention for the orientations. Let $(M,\mathcal{T})$ be a closed aspherical framed 3-manifold endowed with a torus splitting and denote by $p\colon\ (\widetilde{M},\widetilde{\mathcal{T}})\to(M,\mathcal{T})$ a finite covering endowed with a torus splitting defined by $\widetilde{\mathcal{T}}=p^{-1}(\mathcal{T})$. Let (Σ,h) denote a Seifert piece of $M\setminus\mathcal{T}$, where h denotes the fiber of Σ and let $(\Sigma_1,h_1),...,(\Sigma_l,h_l)$ the components of $p^{-1}(\Sigma)$ where h_i denotes the fiber of Σ_i so that $p|\Sigma_i\colon\ \Sigma_i\to\Sigma$ is a fiber preserving map (such a Seifert fibration always exists on Σ_i by [JS]). Then we orient the fibers h_i so that $p|h_i\colon\ h_i\to h$ is orientation preserving. In this case we say that $\widetilde{M}$ is endowed with the framing induced by that of M.

Proposition 2.4. *Let $(M, \mathcal{T})$ be a closed aspherical framed 3-manifold endowed with a torus splitting. Any finite covering $p\colon \widetilde{M} \to M$ induces isometries $p_\sharp\colon H_3\left(\widetilde{M}; \mathbf{R}\right) \to H_3(M; \mathbf{R})$ and*

$$p_\sharp | \widehat{H}_2^{l_1,+}\left(\widetilde{M}^{\widetilde{\mathcal{T}}}; \mathbf{R}\right) : \widehat{H}_2^{l_1,+}\left(\widetilde{M}^{\widetilde{\mathcal{T}}}; \mathbf{R}\right) \to \widehat{H}_2^{l_1}(M; \mathbf{R})$$

where $\widetilde{\mathcal{T}}$ is the torus splitting of $\widetilde{M}$ equal to $p^{-1}(\mathcal{T})$ and where $\widetilde{M}$ is endowed with the framing induced by that of M.

Note that the above *covering property* gives rise to the following

Question 1. Let M be a closed orientable aspherical 3-manifold and let $p\colon \widetilde{M} \to M$ denote a Haken finite covering of M.

(i) Does the covering induce an isometry $p_\sharp | \widehat{H}_2^{l_1,+}\left(\widetilde{M}^{\widetilde{\mathcal{T}}}; \mathbf{R}\right) : \widehat{H}_2^{l_1,+}\left(\widetilde{M}^{\widetilde{\mathcal{T}}}; \mathbf{R}\right) \to \widehat{H}_2^{l_1}(M; \mathbf{R})$ for any torus splitting $\widetilde{\mathcal{T}}$ of $\widetilde{M}$?

(ii) More generally does the covering induce an isometry $p_\sharp\colon \widehat{H}_2^{l_1}\left(\widetilde{M}; \mathbf{R}\right) \to \widehat{H}_2^{l_1}(M; \mathbf{R})$?

2.2. **The secondary fundamental class of a manifold.** We can now define the secondary fundamental class of a closed orientable aspherical 3-manifold M. Consider a torus splitting $\mathcal{T}$ of M and denote by $P_1, ..., P_k$ the components of $M \setminus \mathcal{T}$ supporting a Seifert fibration. Then we set

$$\|M\|^{\mathcal{T}} = \|\widehat{\Theta}_{P_1}(\alpha_{P_1}) + ... + \widehat{\Theta}_{P_k}(\alpha_{P_k})\|_1$$

The class $\widehat{\Theta}_{P_1}(\alpha_{P_1}) + ... + \widehat{\Theta}_{P_k}(\alpha_{P_k}) \in \widehat{H}_2^{l_1,+}\left(M^{\mathcal{T}}; \mathbf{R}\right)$ will be denoted by $[M]^{\mathcal{T}}$ and will be termed the secondary fundamental class of $(M, \mathcal{T})$.

Question 2. Is it possible to compare $[M]^{\mathcal{T}_1}$ and $[M]^{\mathcal{T}_2}$ in $\widehat{H}_2^{l_1}(M; \mathbf{R})$ when $\mathcal{T}_1$ and $\mathcal{T}_2$ are two distinct torus splittings of M?

2.3. **Organization of the paper.** In section 3 we prove Lemmas 2.1 and 2.2 and we recall some results related to the duality between bounded cohomology and l_1-homology. Section 4 will be devoted the construction of bounded 2-cocyles which roughly speaking measure the area of the horizontal surfaces passing through the Seifert pieces of a manifold. This kind of cocyles will be used the estimate the l_1-norm certain l_1-homology classes (see Proposition 4.1). In section 5 we prove Theorem 2.3 and Proposition 2.4 and section 6 is devoted to the proof of Theorem 1.1 and Corollary 1.2.

3. FILLING HOMOMORPHISMS AND DUALITY BETWEEN BOUNDED COHOMOLOGY AND l_1-HOMOLOGY

Proof of Lemma 2.1. Let $\alpha \in H_2(Y, A)$ and choose a relative 2-cycle z in (Y, A) which represents α. Then $\partial z \in Z_1(A)$ and since $H_1^{l_1}(A) = 0$ then there exists $u \in C_2^{l_1}(A)$ such that $\partial u = -\partial z$ in such a way that $z + u \in Z_2^{l_1}(Y)$. This proves point (i).

On the other hand the class of $z + u$ does not depend on the choice of u. Indeed let u_1, u_2 be elements of $C_2^{l_1}(A)$ such that $\partial u_1 = \partial u_2 = -\partial z$. Then $(z + u_1) - (z + u_2) = u_1 - u_2 \in Z_2^{l_1}(A)$. Then there exists $w \in C_3^{l_1}(A)$ such that $\partial w = (z + u_1) - (z + u_2)$.

Moreover the class of $z + u$ does not depend on the choice of the representant z. Indeed, let z and z' be two relative 2-cycles which represent α. Then there exists $p \in C_2(A)$ and $v \in C_3(Y)$ such that $z - z' = \partial v + p$. Let u and u' in $C_2^{l_1}(A)$ such that $\partial u = -\partial z$ and

$\partial u' = -\partial z'$. Then $(z+u)-(z'+u') = \partial v + (p+u-u')$. Thus $p+u-u' \in Z_2^{l_1}(A)$. Then there exists $w \in C_3^{l_1}(A)$ such that $\partial w = p+u-u'$. Thus the map $z \mapsto z+u$ gives rise to a homomorphism

$$\Theta_Y\colon\ H_2(Y,A) \to H_2^{l_1}(Y) \to H_2^{l_1}(X)$$

defined by $\Theta_Y([z]) = [z+u]$. This proves point (ii).

Let $\alpha \in H_2(Y,A)$ and fix $\varepsilon > 0$. Then by [G, Equivalence Theorem] there exists a representative z of α such that $\|z\|_1 \leq \|\alpha\|_1+\varepsilon$ and $\|\partial z\|_1 \leq \varepsilon$. By the Uniform Boundary Condition (see [MM, Theorem 2.3]) there exists a constant $K > 0$ which only depends on the dimension and $u \in C_2^{l_1}(A)$ such that $\partial u = -\partial z$ and $\|u\|_1 \leq K\|\partial z\|_1 < K\varepsilon$. This implies, passing to the limits, that $\|\Theta_Y(\alpha)\|_1 \leq \|\alpha\|_1$. This proves point (iii).

Let α be an element of $H_2(Y,A;\mathbf{R})$. Then $f_\sharp(\Theta_Y\alpha) = [f_\sharp(z) + f_\sharp(u)]$ where z is a relative 2-cycle representing α and u is a l_1 2-chain in A such that $\partial u = -\partial z$. Since $\partial f_\sharp u = -\partial f_\sharp z$ and since $f_\sharp u$ is a l_1 2-chain in B then $[f_\sharp(z)+f_\sharp(u)] = \Theta_W(f_\sharp(\alpha))$. This proves of point (iv).

□

Proof of Lemma 2.2. Let (P,h) be a Seifert oriented 3-manifold with a fixed fibration h. Let $p\colon\ \widetilde{P} \to P$ denote a finite covering of P. Then p induces an epimorphism $H_2(\widetilde{P},\partial\widetilde{P}) \to H_2(P,\partial P)$ and thus passing to the quotient we get an epimorphism $\widehat{H}_2(\widetilde{P},\partial\widetilde{P}) \to \widehat{H}_2(P,\partial P)$.

If P admit a geometry Nil, R^3 or $\widetilde{\mathrm{SL}}(2,\mathbf{R})$ then it admits a finite covering $\widetilde{P}$ which is either a torus bundle (over $\mathbf{S}^1$ or I) or a circle bundle over a hyperbolic surface with non-zero Euler number. In any case $\widehat{H}_2(\widetilde{P},\partial\widetilde{P}) = \{0\}$. Indeed in these cases $\widetilde{P}$ contains no incompressible surfaces with negative Euler characteristic. This proves that $\widehat{H}_2(P,\partial P)$ is trivial.

Hence from now on we assume that $\mathrm{int}(P)$ admits a $\mathbf{H}^2\times\mathbf{R}$-geometry. Let $p\colon\ \widetilde{P} \to P$ denotes a finite regular covering of P homeomorphic to a product $\widetilde{F}\times\mathbf{S}^1$. Note that since $\widetilde{P}$ has an orientable base then by [WZ, Lemma 6] $H_2(\widetilde{P},\partial\widetilde{P})$ is generated by any minimal surface together with vertical surfaces (which are either tori or vertical annuli) and thus $\widehat{H}_2(\widetilde{P},\partial\widetilde{P})$ is generated by any minimal surface. This proves that $\widehat{H}_2(\widetilde{P},\partial\widetilde{P}) = \mathbf{R}$ and thus $\widehat{H}_2(P,\partial P) = \mathbf{R}$.

Let us check the second statement of the lemma. Let F_0 and F_1 denote two minimal surfaces in P and denote by $\widetilde{F}_0$, resp. $\widetilde{F}_1$ the spaces $p^{-1}(F_0)$, resp. $p^{-1}(F_1)$. Since F_0 and F_1 are minimal then in particular $\chi(F_0) = \chi(F_1)$ and thus $\chi(\widetilde{F}_0) = \chi(\widetilde{F}_1)$ which implies, since $\widetilde{P}$ is a product that $[\widetilde{F}_0,\partial\widetilde{F}_0] = [\widetilde{F}_1,\partial\widetilde{F}_1]$ in $\widehat{H}_2(\widetilde{P},\partial\widetilde{P})$. Since for each $i=0,1$ we have $p_\sharp([\widetilde{F}_i,\partial\widetilde{F}_i]) = \deg(p)[F_i,\partial F_i]$ then $[F_0,\partial F_0] = [F_1,\partial F_1]$ which proves the lemma.

□

Throughout this paper we will need the following general results which come from the duality between l_1-chains and bounded cochains.

Lemma 3.1. *The Kronecker product between l_1-homology and bounded cohomology gives rise to a surjective bounded operator*

$$\Phi\colon\ \widehat{H}_b^*(X) \to (\widehat{H}_*^{l_1}(X))'$$

where $(\widehat{H}^{l_1}_*(X))'$ *denotes the space of continuous linear forms on* $\widehat{H}^{l_1}_*(X)$*, such that* $\|\Phi\| = 1$ *and for any* $\varphi \in (\widehat{H}^{l_1}_*(X))'$ *there exists* $\beta \in \widehat{H}^*_b(X)$ *such that* $\Phi(\beta) = \varphi$ *and* $\|\beta\|_\infty = \|\varphi\|_\infty$*.*

Proof. This follows directly from the Holder inequality combined with the Hahn-Banach Theorem. ☐

Let Γ be a group acting by homeomorphisms on a topological space X. We denote by $\Gamma\widehat{H}^{l_1}_*(X) = \{\alpha \in \widehat{H}^{l_1}_*(X) \text{ s.t. } g_\sharp(\alpha) = \alpha \text{ when } g \in \Gamma\}$.

Lemma 3.2. *Let* $p\colon\ \widetilde{X} \to X$ *be a regular covering map with finite Galois group* Γ*. Then the induced homomorphism* $p_\sharp\colon\ \Gamma\widehat{H}^{l_1}_*(\widetilde{X}) \to \widehat{H}^{l_1}_*(X)$ *is an isometry.*

Proof. Consider the averaging retraction $A\colon\ C^*_b(\widetilde{X}) \to C^*_b(X)$ defined by

$$\langle A(\gamma), \sigma\rangle = \frac{\sum_{g\in\Gamma}\langle g^*\gamma, \widetilde{\sigma}\rangle}{\mathrm{Card}(\Gamma)}$$

where $\widetilde{\sigma}\colon\ \Delta^* \to \widetilde{X}$ denotes a lifting of $\sigma\colon\ \Delta^* \to X$. This definition does not depend one the choice of the lifting $\widetilde{\sigma}$ since the covering is regular. By construction A satisfies the identity $A\circ p^* = \mathrm{Id}$ and commutes with the differentials so that it induces a homomorphism $\widehat{A}\colon\ H^*_b(\widetilde{X}) \to H^*_b(X)$. Then at the H^*_b-level we still have the identity $\widehat{A}\circ p^* = \mathrm{id}$ and $\|\widehat{A}\| \leq 1$. Notice that $\widehat{A}$ induces a homomorphism $\widehat{H}^*_b(\widetilde{X}) \to \widehat{H}^*_b(X)$ still denoted by $\widehat{A}$.

Let $\alpha = \overline{[z]} \in \Gamma\widehat{H}^{l_1}_*(\widetilde{X})$, where $\overline{-}\colon\ H^{l_1}_*(\widetilde{X}) \to \widehat{H}^{l_1}_*(\widetilde{X})$ denotes the natural quotient homomorphism and z is a l_1-cycle. If $\alpha = \overline{[z]} \neq 0$ then by Lemma 3.1, there exists $\beta = \overline{[\gamma]} \in \widehat{H}^*_b(\widetilde{X})$ such that $\langle\beta, \alpha\rangle = 1$ and $\|\beta\|_\infty = \frac{1}{\|\alpha\|_1}$. Since α is Γ-invariant then by the definition of the averaging we have the equalities

$$\left\langle \widehat{A}(\beta), p_\sharp(\alpha)\right\rangle = \langle A(\gamma), p_\sharp(z)\rangle = \frac{1}{\mathrm{Card}(\Gamma)}\sum_{g\in\Gamma}\langle g^*(\gamma), z\rangle = \langle\gamma, z\rangle = \langle\beta, \alpha\rangle = 1$$

and thus

$$\|p_\sharp(\alpha)\|_1 \geq \frac{1}{\|\widehat{A}(\beta)\|_\infty} \geq \frac{1}{\|\beta\|_\infty} = \|\alpha\|_1$$

Since the inequality $\|p_\sharp\| \leq 1$ is always true this proves the lemma. ☐

4. BOUNDED 2-COCYLES MEASURING THE HORIZONTAL AREAS

Let M be an orientable closed aspherical 3-manifold endowed with a torus splitting $\mathcal{T}$. A Seifert piece P of $(M, \mathcal{T})$ will be termed a *product component* if it is homeomorphic to a product $F \times \mathbf{S}^1$, where F is a surface whose interior admits a hyperbolic structure. The purpose of this section is to construct for each "product component" P of $M \setminus \mathcal{T}$, a bounded 2-cocyle in M which measures the hyperbolic area of horizontal surfaces of P. More precisely the main result of this section states as follows:

Proposition 4.1. *For each product component* $P = F \times \mathbf{S}^1$ *of* $M \setminus \mathcal{T}$ *there exists a non-trivial bounded* 2*-cocyle* Ω_P *in* M *satisfying the following properties:*

(i) $i^*(\Omega_P)$ *is a relative* 2*-cocycle in* $(P, \partial P)$ *where* $i\colon\ P \hookrightarrow M$ *denotes the natural inclusion,*

(ii) for any connected horizontal surface $\mathcal{F}$ *in* P *then*

$$\left|\left\langle [\Omega_P], \widehat{\Theta}_P([\mathcal{F}])\right\rangle\right| = |\langle i^*(\Omega_P), z_{\mathcal{F}}\rangle| = \mathrm{Area}(\mathcal{F})$$

where $z_{\mathcal{F}}$ is a relative 2-cycle representing the fundamental class of $\mathcal{F}$ and $\mathrm{Area}(\mathcal{F})$ denotes the area of $\mathrm{int}(\mathcal{F})$ endowed with a complete hyperbolic metric.

Let $(P_i)_{i\in I}$ be a family of pairwise distinct product components of $M \setminus \mathcal{T}$ and for each $i \in I$ denote by $k_i\colon\ P_i \to M$ the canonical inclusion. Then

(iii) $k_i^*(\Omega_{P_j}) = 0$ if $i \neq j$ and

$$\left\| \sum_{i\in I} [\Omega_{P_i}] \right\|_\infty = \pi$$

where $[\Omega_{P_i}]$ denotes the class of Ω_{P_i} in $\widehat{H}_b^2(M;\mathbf{R})$ for each $i \in I$.

To prove this proposition, we first need to construct a chain map: the *reduction*, used in [FS] (see section 4.1). Next we *straight horizontally* the reduced chains which meet essentially P to define a bounded 2-cochain Ω_P (see section 4.2). The proof of Proposition 4.1 will occupy section 4.3. From now on, we denote by Δ^n the standard n-simplex defined by

$$\Delta^n = [v_0,...,v_n] = \left\{ \sum_{i=0}^n t_i v_i, \sum_{i=0}^n t_i = 1, t_i \geq 0 \right\}$$

where $v_i = (0,...,1,...0) \in \mathbf{R}^{n+1}$. We denote by $V(\Delta^n) = \{v_0,...,v_n\}$ the vertices of Δ^n and by Δ_i^{n-1} the i-th $(n-1)$-face of Δ^n defined by $\Delta_i^{n-1} = [v_0,...,\widehat{v_i},...,v_n]$. Note that Δ^n is oriented by the order of its vertices in such a way that $(v_1 - v_0,...,v_n - v_0)$ is a positive frame.

4.1. **Reduction operator in singular homology.**

Let M be a closed aspherical orientable 3-manifold endowed with an amenable splitting $\mathcal{T}$. Denote by $P_1,...,P_l$ the components of $M \setminus \mathcal{T}$. As in [FS], we consider a chain map $\rho\colon\ C_n(M) \to C_n(M)$ defined as follows:

If $n = 0$ then we set $\rho = \mathbf{1}$.

If $n = 1$ let $\tau\colon\ [v_0,v_1] \to M$ be a 1-simplex. Since $\mathcal{T}$ is incompressible, the map τ is homotopic rel. $\{v_0,v_1\}$ to a *reduced* 1-simplex i.e. a map $\tau_1\colon\ [v_0,v_1] \to M$ such that either $\tau_1([v_0,v_1]) \subset \mathcal{T}$ or $\tau_1|(v_0,v_1)$ is transverse to $\mathcal{T}$ and for each component J of $\tau_1^{-1}(P_i)$ then $\tau_1|J$ is not homotopic rel. ∂J into ∂P_i. Then we set $\rho(\tau) = \tau_1$ and we extend ρ by linearity.

If $n = 2$ let $\sigma\colon\ \Delta^2 = [v_0,v_1,v_2] \to M$ be a 2-simplex. Then σ is homotopic rel. $V(\Delta^2) = \{v_0,v_1,v_2\}$ to a reduced 2-simplex σ_1 such that either $\sigma_1(\Delta^2) \subset \mathcal{T}$ or $\sigma_1|\mathrm{int}(\Delta^2)$ is transverse to $\mathcal{T}$, the 1-simplex $\sigma_1|e$ is reduced for each edge e of Δ^2 and $\sigma_1^{-1}(\mathcal{T})$ contains no loop components. Thus if J is a component of $\sigma_1^{-1}(\mathcal{T})$ such that $J \cap \mathrm{int}(\Delta^2) \neq \emptyset$ then J is a proper arc in Δ^2 connecting two distinct edges. Then we set $\rho(\sigma) = \sigma_1$ and we extend ρ by linearity.

Remark 4.2. Let $\sigma\colon\ \Delta^2 \to M$ be a reduced 2-simplex. Denote by D a component of $\sigma^{-1}(\mathrm{int}(P))$, where P is a component of $M \setminus \mathcal{T}$. Each component of $V(\Delta^2) \cap D$ will be termed a vertex of D and each component of $\sigma^{-1}(\partial P) \cap \overline{D}$ will be termed an *ideal vertex* of D. Thus an ideal vertex of D is either a vertex or an edge or a proper arc of Δ^2 connecting two distinct edges. Let us denote by $V_\infty(D)$ the set of ideal vertices of D and by $V(D)$ the set of all vertices and ideal vertices of D. Note that if $D \neq \emptyset$ then $2 \leq \mathrm{Card}(V(D)) \leq 3$.

Remark 4.3. Suppose that $\sigma\colon\ \Delta^2 \to M$ is a reduced 2-simplex. If $\sigma(e)$ is not contained in $\mathcal{T}$ for any edge e of Δ^2 then there exists a unique component, denoted by $\mathrm{Core}(\sigma)$, of $\Delta^2 \setminus \sigma^{-1}(\mathcal{T})$ which meets the three edges of Δ^2 (see [FS]). On the other hand, if e_1 and

e_2 denote two distinct edges of Δ^2 such that $\sigma(e_1) \subset \mathcal{T}$ and $\sigma(e_2) \subset \mathcal{T}$ then $\sigma(\Delta^2) \subset \mathcal{T}$ (this follows directly from the reduction hypothesis of σ).

If $n = 3$ let $\sigma\colon\ \Delta^3 = [v_0, v_1, v_2, v_3] \to M$ be a 3-simplex. Then σ is homotopic rel. $V(\Delta^3) = \{v_0, v_1, v_2, v_3\}$ to a reduced 3-simplex σ_1 such that either $\sigma_1(\Delta^3) \subset \mathcal{T}$ or $\sigma_1|\mathrm{int}(\Delta^3)$ is transverse to $\mathcal{T}$, the 2-simplex $\sigma_1|\Delta^2_i$ is reduced for each face Δ^2_i of Δ^3 and if D is a component of $\sigma_1^{-1}(\mathcal{T})$ such that $D \cap \mathrm{int}(\Delta^3) \neq \emptyset$ then D is either a normal triangle or a normal rectangle (see figure 1).

Then we set $\rho(\sigma) = \sigma_1$ and we extend ρ by linearity. For $n = 0, 1, 2, 3$ we denote by $C_n^{\mathrm{red}}(M)$ the image of $C_n(M)$ under ρ. Notice that the reduction is stable under the ∂-operator.

Remark 4.4. Let σ denote a reduced 3-simplex and let ∇ be a component of $\sigma^{-1}(\mathrm{int}(P))$, where P is a component of $M \setminus \mathcal{T}$. Each component of $V(\Delta^3) \cap \nabla$ will be termed a vertex of ∇ and each component of $\sigma^{-1}(\partial P) \cap \overline{\nabla}$ which is a normal triangle or a vertex of Δ^3 will be termed an *ideal vertex* of ∇. Denote by $V_\infty(\nabla)$ the ideal vertices of ∇ and by $V(\nabla)$ all the vertices and ideal vertices of ∇. The set $\sigma^{-1}(\partial P) \cap \overline{\nabla}$ consists of either

(i) exactly one 2-face Δ^2_i of Δ^3 and at most one ideal vertex of ∇^3, or

(ii) exactly two edges of Δ^3 with no common vertices, or

(iii) exactly one edge and at most one normal rectangle or at most two ideal vertices, or

(iv) no faces and no edges components and at most four ideal vertices or two rectangles or one rectangle and at most two ideal vertices.



Figure 1: Normal triangles and rectangles

4.2. **Horizontal area.** Suppose now that M is endowed with a torus splitting $\mathcal{T}$. Let P be a product component of $M \setminus \mathcal{T}$ identified with a product $F \times \mathbf{S^1}$, where F is an orientable surface whose interior W admits a hyperbolic structure. Denote by $q\colon\ P \to F$ the projection to the first factor.

Let $\sigma\colon\ \Delta^2 \to M$ be a reduced 2-simplex and let D be a component of $\sigma^{-1}(\mathrm{int}(P))$. Denote by σ_D the map $\sigma|D\colon\ D \to \mathrm{int}(P)$ and by $\widetilde{q\sigma_D}\colon\ D \to \mathbf{H}^2$ a lifting of $q\sigma_D\colon\ D \to W$ into the universal covering of W. It follows from Remark 4.2 that D has 2 or 3 vertices (including ideal vertices). Denote by $V(D) = \{w_0, w_1, w_2\}$ the set of vertices of D (may be $w_i = w_j$ for some $i \neq j$) in such a way that if $D = \mathrm{Core}(\sigma)$ then each w_i corresponds to v_i (see figure 2).



Figure 2

For each $i = 0, 1, 2$, the vertex w_i has a corresponding image w_i^∞ in $\overline{\mathbf{H}}^2 = \mathbf{H}^2 \cup \partial_\infty \mathbf{H}^2$ defined as follows:

If $w_i \notin V_\infty(D)$ then $w_i^\infty = \widetilde{q\sigma_D}(w_i)$. If $w_i \in V_\infty(D)$ denote by w_j a vertex of D distinct from w_i. Choose a point x_i in w_i and x_j in w_j and consider the open geodesic segment (x_i, x_j) in Δ^2. Then there exists $\varepsilon > 0$ and a horodisk C_i in $\mathbf{H}^2$ such that $\widetilde{q\sigma_D}((x_i, x_i + \varepsilon(x_j - x_i)]) \in C_i$ and in this case w_i^∞ denotes the center of C_i (which is defined as the contact point of C_i with $\partial_\infty \mathbf{H}^2$, see figure 3). Note that w_i^∞ does not depend on the choice of the points x_i and x_j and does not depend on the choice of $w_j \neq w_i$ since in any case it follows from our construction and from the reduction hypothesis on σ that

$$\lim_{t \to 1^-} \widetilde{q\sigma_D}(tx_i + (1-t)x_j) = w_i^\infty$$

for any $x_i \in w_i$, $x_j \in w_j$ and $w_j \neq w_i$. On the other hand, since σ is reduced, if $w_i \neq w_j$ where w_i or w_j is an ideal vertex then $w_i^\infty \neq w_j^\infty$. The set $\{w_0^\infty, w_1^\infty, w_2^\infty\}$ will be termed the σ-image of the (ideal) vertices of D.



(I)

(II)

Figure 3

Let $\{w_0^\infty, w_1^\infty, w_2^\infty\}$ denote three points (note necessarily pairwise distinct) of $\overline{\mathbf{H}}^2$. Then we denote by $[w_0^\infty, w_1^\infty, w_2^\infty]$ the straight 2-simplex of $\overline{\mathbf{H}}^2$ spanned by $\{w_0^\infty, w_1^\infty, w_2^\infty\}$ and oriented by the order of its vertices. Then we associate to each component D of $\sigma^{-1}(\mathrm{int}(P))$ an algebraic area denoted by $\mathcal{A}_{\mathbf{H}}(D)$ defined by $\pm$ the hyperbolic area of $[w_0^\infty, w_1^\infty, w_2^\infty]$ depending on whether the orientation of $[w_0^\infty, w_1^\infty, w_2^\infty]$ matches the orientation of $\mathbf{H}^2$ or not. Notice that the area $\mathcal{A}_{\mathbf{H}}(D)$ does not depend on the chosen lifting $\widetilde{q\sigma_D}\colon\ D \to \mathbf{H}^2$ of $q\sigma_D\colon\ D \to W$ into the universal covering $\mathbf{H}^2$ of W.

Let us define a 2-cochain in M in the following way. Let $\sigma\colon\ \Delta^2 \to M$ denote a singular 2-simplex. If $\rho\sigma(\Delta^2) \subset M \setminus \mathrm{int}(\mathrm{P})$ then we set

$$\langle \Omega_P, \sigma \rangle = 0$$

and if $\rho\sigma(\Delta^2) \cap \mathrm{int}(\mathrm{P}) \neq \emptyset$ then we set

$$\langle \Omega_P, \sigma \rangle = \sum_{D \in \rho(\sigma)^{-1}(\mathrm{int}(P))} \mathcal{A}_{\mathbf{H}}(D)$$

4.3. **Proof of Proposition 4.1.** We first check that Ω_P is a cocycle. Let $\sigma\colon\ \Delta^3 \to M$ be a 3-simplex. Then we have by definition of Ω_P

$$\langle \delta\Omega_P, \sigma\rangle = \langle \delta\Omega_P, \rho(\sigma)\rangle = \sum_{i=0}^{3} (-1)^i \langle \Omega_P, \rho(\sigma)|\Delta_i^2\rangle$$

where Δ_i^2 denotes the i-th face $[v_0, ..., \widehat{v}_i, ..., v_3]$ of Δ^3. To check the cocyle condition one can always assume that $\rho\sigma(\Delta^3) \cap \mathrm{int}(P) \neq \emptyset$. Thus by the definition of Ω_P we have

$$\langle \delta\Omega_P, \sigma\rangle = \sum_{i=0}^{3} (-1)^i \sum_{D \in (\rho\sigma|\Delta_i^2)^{-1}(\mathrm{int}(P))} \mathcal{A}_{\mathbf{H}}(D)$$

Notice that for each component D of $(\rho\sigma|\Delta_i^2)^{-1}(\mathrm{int}(P))$ there exists a component ∇ of $\rho(\sigma)^{-1}(\mathrm{int}(P))$ whose D is a 2-face. Thus to check the cocyle condition it is sufficient to prove that for any component ∇ of $\rho(\sigma)^{-1}(\mathrm{int}(P))$ then

$$\sum_{i=0}^{3} (-1)^i \mathcal{A}_{\mathbf{H}}(\partial_i\nabla) = 0$$

where $\partial_i\nabla = \nabla \cap \Delta_i^2$ for $i = 0, ..., 3$, with the obvious convention $\mathcal{A}_{\mathbf{H}}(\partial_i\nabla) = 0$ if $\sigma(\partial_i\nabla) \subset \partial P$.

1. Assume that $\overline{\nabla} \cap (\rho\sigma)^{-1}(\partial P)$ contains a 2-face Δ_i^2 of Δ^3. Then in this case each $\partial_j\nabla$, for $j \neq i$, has only 2 vertices and thus $\mathcal{A}_{\mathbf{H}}(\partial_i\nabla) = 0$ for $i = 0, ..., 3$.

2. Assume that $\overline{\nabla} \cap (\rho\sigma)^{-1}(\partial P)$ contains no face of Δ^3 and two edges, say e_1 and e_2 of Δ^3. Since by Remark 4.3 we have $\overline{e}_1 \cap \overline{e}_2 = \emptyset$ then as in Case 1 each $\partial_i\nabla$ has only 2 vertices.

3.1 Assume that $(\rho\sigma)^{-1}(\partial P) \cap \overline{\nabla}$ contain no face, one edge, say $[v_i, v_j]$ with $i < j$ of Δ^3 and a normal rectangle R. Then again in this case each $\partial_i\nabla$ has only 2 vertices.

3.2. Assume that $(\rho\sigma)^{-1}(\partial P) \cap \overline{\nabla}$ contain no face, one edge $[v_i, v_j]$ of Δ^3 and at most two ideal vertices corresponding to v_k and v_l.

If $i < j < k < l$ then $\mathcal{A}_{\mathbf{H}}(\partial_\nu\nabla) = 0$ when $\nu = k, l$ since in this case $\partial_\nu\nabla$ has only 2 vertices. Denote by $\{w_i, w_k, w_l\}$, resp. $\{w_j, w_k, w_l\}$, the vertices of $\partial_j\nabla$, resp. of $\partial_i\nabla$. Choose a base point x in the open edge (w_k, w_l) and a corresponding base point $\widetilde{x}$ in $\mathbf{H}^2$ over $q\rho\sigma(x)$. Denote by $\{w_i^\infty, w_k^\infty, w_l^\infty\}$, resp. $\{w_j^\infty, w_k^\infty, w_l^\infty\}$, the σ-images of $\{w_i, w_k, w_l\}$, resp. $\{w_j, w_k, w_l\}$ corresponding to the lifting $\widetilde{q\rho\sigma}_{|\partial_j\nabla}$ of $q\rho\sigma_{|\partial_j\nabla}$, resp. $\widetilde{q\rho\sigma}_{|\partial_i\nabla}$ of $q\rho\sigma_{|\partial_i\nabla}$, such that $\widetilde{q\rho\sigma}_{|\partial_j\nabla}(x) = \widetilde{q\rho\sigma}_{|\partial_i\nabla}(x) = \widetilde{x}$. Notice that since $\rho\sigma([v_i, v_j]) \subset \partial P$ then $w_i^\infty = w_j^\infty$. Thus in this case, since $j = i + 1$, we have the equality

$$(-1)^j \mathcal{A}_{\mathbf{H}}(\partial_j\nabla) + (-1)^i \mathcal{A}_{\mathbf{H}}(\partial_i\nabla) = 0$$

If $i < k < j < l$ then $\mathcal{A}_{\mathbf{H}}(\partial_\nu\nabla) = 0$ when $\nu = k, l$. Denote by $\{w_i, w_k, w_l\}$, resp. $\{w_k, w_j, w_l\}$, the vertices of $\partial_j\nabla$, resp. of $\partial_i\nabla$. Denote by $\{w_i^\infty, w_k^\infty, w_l^\infty\}$, resp. $\{w_k^\infty, w_j^\infty, w_l^\infty\}$, the σ-images of $\{w_i, w_k, w_l\}$, resp. $\{w_k, w_j, w_l\}$. Since $\rho\sigma([v_i, v_j]) \subset \partial P$ then $w_i^\infty = w_j^\infty$. Thus in this case we have the equality

$$(-1)^j \mathcal{A}_{\mathbf{H}}(\partial_j\nabla) + (-1)^i \mathcal{A}_{\mathbf{H}}(\partial_i\nabla) = 0$$

since $j = i + 2$ and $[w_i^\infty, w_k^\infty, w_l^\infty]$ and $[w_k^\infty, w_j^\infty, w_l^\infty]$ are the same geodesic simplices with opposite orientations. The other possibilities for i, j, k, l follow in the same way.

Assume now that $(\rho\sigma)^{-1}(\partial P) \cap \overline{\nabla}$ consists only of vertices, normal triangles or normal rectangles of Δ^3. Note that it follows from the construction that the only possibilities are $\mathrm{Card}(V(\nabla)) = 0, 2, 4$.

4.1. Assume $\mathrm{Card}(V(\nabla)) = 4$ and denote by w_0, w_1, w_2, w_3 the vertices of ∇ corresponding to v_0, v_1, v_2, v_3 and denote by $w_0^\infty, w_1^\infty, w_2^\infty, w_3^\infty$ their σ-images. Then

$$\sum_{i=0}^{3}(-1)^i[w_0^\infty, ..., \widehat{w_i^\infty}, ..., w_3^\infty]$$

is the boundary of a geodesic 3-simplex in $\mathbf{H}^2$. Using the same argument as in [BG] we know that there are two distinct configurations for a geodesic 3-simplex in $\mathbf{H}^2$ and in any case we have

$$\sum_{i=0}^{3}(-1)^i \mathcal{A}_{\mathbf{H}}(\partial_i \nabla) = 0$$

4.2. Assume $\mathrm{Card}(V(\nabla)) = 2$. Denote by w_i and w_j the vertices of ∇. Then we have the two following possibilities:

4.2.1. Suppose first that $\partial\nabla$ contains a normal rectangle R. We denote by v_i, v_j the vertices of Δ^3 corresponding to the vertices w_i and w_j of ∇. First notice that $\partial_i\nabla$ and $\partial_j\nabla$ contain only two vertices: w_j, resp. w_i, and the vertex corresponding to $R \cap \partial_i\nabla$, resp. $R \cap \partial_j\nabla$. Then

$$\mathcal{A}_{\mathbf{H}}(\partial_i\nabla) = \mathcal{A}_{\mathbf{H}}(\partial_j\nabla) = 0$$

On the other hand $\partial_k\nabla$ and $\partial_l\nabla$ have three vertices: the vertices w_i and w_j and one ideal vertex (denoted by w_l, resp. w_k) "corresponding" to R. Since w_l and w_k correspond to edges of the same rectangle then $w_k^\infty = w_l^\infty$ (for some lifting with the same base point). Then we get again as in 3.2

$$(-1)^k \mathcal{A}_{\mathbf{H}}(\partial_k\nabla) + (-1)^l \mathcal{A}_{\mathbf{H}}(\partial_l\nabla) = 0$$

4.2.2 Suppose that $\partial\nabla$ contains no normal rectangle R then

either $V(\nabla)$ consists of one normal triangle and one vertex of Δ^3. Then $\mathcal{A}_{\mathbf{H}}(\partial_i\nabla) = 0$ for any i, since $\partial_i\nabla$ contains only two vertices.

either $V(\nabla)$ consists of two parallel normal triangles. Then again $\mathcal{A}_{\mathbf{H}}(\partial_i\nabla) = 0$ for any i, since $\partial_i\nabla$ contains only two vertices.

4.3. Assume $\mathrm{Card}(V(\nabla)) = 0$. Then necessarily, $\partial\nabla$ contains two parallel rectangles and $\mathcal{A}_{\mathbf{H}}(\partial_i\nabla) = 0$ for any i since $\partial_i\nabla$ contains only two vertices. This completes the proof that Ω_P is a cocyle.

Next point (i) of the proposition follows directly from the construction of Ω_P. On the other hand given $\alpha \in \widehat{H}_2(P, \partial P; \mathbf{R})$ and z_α a relative 2-cycle representing of α, we get the following equalities

$$\left\langle [\Omega_P], \widehat{\Theta}_P(\alpha) \right\rangle = \langle \Omega_P, z_\alpha + u_\alpha \rangle = \langle i^*(\Omega_P), z_\alpha \rangle = \langle [i^*(\Omega_P)], [z_\alpha] \rangle$$

where u_α is a l_1-chain in ∂P such that $\partial u_\alpha = -\partial z_\alpha$. In particular, if z_α represents the fundamental class of F then we get

$$\left\langle [\Omega_P], \widehat{\Theta}_P(\alpha) \right\rangle = \mathrm{Area}(F)$$

Indeed we can choose as a representant z_α the formal sum of a triangulation of F and then apply Ω_P. On the other hand, if $\mathcal{F}$ denotes any connected horizontal surface in P then since $\mathcal{F}$ is a finite covering of F we get $\left\langle [\Omega_P], \widehat{\Theta}_P([\mathcal{F}]) \right\rangle = \mathrm{Area}(\mathcal{F})$. This proves point (ii).

It remains to check point (iii). The fact that $k_i^*(\Omega_{P_j}) = 0$ for any $i \neq j$ follows from the construction of Ω_{P_j}. We first check that $\| \sum_{i \in I} \Omega_{P_i} \|_\infty \leq \pi$. For each $i \in I$ we can identify P_i with the product $F_i \times \mathbf{S}^1$. Let $\sigma\colon \Delta^2 \to M$ be a singular 2-simplex. If there exists an edge e of Δ^2 such that $\rho\sigma(e) \subset \mathcal{T}$ then $\langle \sum_{i\in I} \Omega_{P_i}, \sigma\rangle = 0$. If for any edge e of Δ^2 we have $\rho\sigma(e) \not\subset \mathcal{T}$ then we know from Remark 4.3 that there exists a unique component $\mathrm{Core}(\sigma)$ of $(\rho\sigma)^{-1}(M \setminus \mathcal{T})$ which meets the three edges of Δ^2. Denote by P_ν the component of $M \setminus \mathcal{T}$ such that $\rho\sigma(\mathrm{Core}(\sigma)) \subset \mathrm{int}(P_\nu)$. If D is a component of $(\rho\sigma)^{-1}(\mathrm{int}(P_i))$ for some $i \in I$ distinct from $\mathrm{Core}(\sigma)$ then we get the equality

$$\mathcal{A}_{\mathbf{H}}(D) = 0$$

Indeed in this case D has 2 (ideal) vertices. On the other hand, if $P_\nu = P_{i_0}$ for some $i_0 \in I$ then we know that

$$|\mathcal{A}_{\mathbf{H}}(D)| \leq \pi$$

by the definition of $\mathcal{A}_{\mathbf{H}}(D)$ since the areas of geodesic triangles in $\mathbf{H}^2$ are bounded by π. This proves that $\| \sum_{i\in I} \Omega_{P_i}\|_\infty \leq \pi$.

Denote by Ω the sum $\sum_{i\in I} \Omega_{P_i}$ and let $i \in I$ be a fixed index. Since Θ_{P_i} is a contraction, then using point (ii) we get the following equalities

$$\left|\left\langle [\Omega], \widehat{\Theta}_{P_i}([F_i])\right\rangle\right| = \mathrm{Area}(F_i) \leq \|[\Omega]\|_\infty \left\|\widehat{\Theta}_{P_i}([F_i])\right\|_1 \leq \|[\Omega]\|_\infty \, \|[F_i]\|_1$$

Since $\|[F_i]\|_1 \leq \|F_i\|$ this completes the proof of the proposition since $\mathrm{Area}(F_i) = \pi\|F_i\|$ by [G, Section 0.4] and [Th].

5. A FILLING ISOMETRY FOR RELATIVE CLASSES

This section is devoted to the proof of Theorem 2.3 and of Proposition 2.4. We begin with a special case.

Lemma 5.1. *Theorem 2.3 is true when each Seifert piece of* $(M, \mathcal{T})$ *is homeomorphic to a product* $F \times \mathbf{S}^1$.

Proof. We first check point (i). Let P be a Seifert piece of $M\setminus\mathcal{T}$. Then P is homeomorphic to a product $F \times \mathbf{S}^1$ where F is a hyperbolic orientable surface. By Proposition 4.1 we deduce, using the Holder inequality that $\|\widehat{\Theta}_P([F])\|_1 \geq \|F\| \geq \|[F]\|_1$ and thus finally we get from above $\|\widehat{\Theta}_P([F])\|_1 = \|[F]\|_1 = \|F\|$. Let $\alpha \in \widehat{H}_2(P, \partial P; \mathbf{R})$. Then by Lemma 2.2, there exists $\xi \in \mathbf{R}$ such that $\alpha = \xi\alpha_P$. Then using Proposition 4.1 we get

$$\left|\left\langle [\Omega_P], \widehat{\Theta}_P(\alpha)\right\rangle\right| = |\xi|\mathrm{Area}(F) \leq \pi\|\widehat{\Theta}_P(\alpha)\|_1$$

Hence $\|\widehat{\Theta}_P(\alpha)\|_1 \geq |\xi|\|F\| \geq \|\alpha\|_1$ and thus $\widehat{\Theta}_P$ is an isometry. If α is the class of horizontal surface $\mathcal{F}$ then F is finitely covered by $\mathcal{F}$ and thus ξ is necessarily and integer satisfying $\|\mathcal{F}\| = |\xi|\|F\|$ (actually ξ is the degree of the map induced by the Seifert projection $\mathcal{F} \to F$). This proves point (i).

We check point (ii). Let $P_1, ..., P_l$ denote the Seifert pieces of $M\setminus\mathcal{T}$. For each P_i denote by F_i a minimal surface. Let $(\alpha_1, ...\alpha_l)$ be a l-uple of $\widehat{H}_2(P_1, \partial P_1) \times ... \times \widehat{H}_2(P_l, \partial P_l)$ and denote by $\xi_1, ..., \xi_l$ the real numbers such that for each $i = 1, ..., l$ we have $\alpha_i = \xi_i\alpha_{P_i}$, where α_{P_i} denotes the class of F_i in $\widehat{H}_2^+(P_i, \partial P_i)$. By Proposition 4.1 there exists for each $i \in \{1, ..., l\}$ a bounded 2-cocycle Ω_i in M such that

$$\left\langle [\Omega_i], \widehat{\Theta}_{P_j}\alpha_j \right\rangle = \delta_{ij}|\xi_i|\mathrm{Area}(F_i)$$

for any i, j in $\{1, ..., l\}$. Thus we get

$$\left\langle \sum_i [\Omega_i], \sum_j \widehat{\Theta}_{P_j}(\alpha_j) \right\rangle = \sum_i |\xi_i| \mathrm{Area}(F_i) \leq \pi \left\| \sum_j \widehat{\Theta}_{P_j}(\alpha_j) \right\|_1$$

thus we get

$$\left\| \sum_i \widehat{\Theta}_{P_i}(\alpha_i) \right\|_1 \geq \sum_i |\xi_i| \|F_i\| \geq \sum_i \|\xi_i [F_i]\|_1 = \sum_i \left\| \widehat{\Theta}_{P_i}(\alpha_i) \right\|_1$$

Since the opposite inequality is also true this proves the lemma. □

We now establish the following result which is a special case of Proposition 2.4.

Lemma 5.2. *Let $(M, \mathcal{T})$ be a closed aspherical framed 3-manifold endowed with a torus splitting and let $p\colon \widetilde{M} \to M$ denote a finite regular covering such that each Seifert component of $\widetilde{M} \setminus \widetilde{\mathcal{T}}$ is homeomorphic to a product, where $\widetilde{\mathcal{T}} = p^{-1}(\mathcal{T})$. Then the covering p induces an isometry $p_\sharp | \widehat{H}_2^{l_1,+}(\widetilde{M}^{\widetilde{\mathcal{T}}}; \mathbf{R})\colon \widehat{H}_2^{l_1,+}(\widetilde{M}^{\widetilde{\mathcal{T}}}; \mathbf{R}) \to \widehat{H}_2^{l_1}(M; \mathbf{R})$ where $\widetilde{M}$ is endowed with the framing induces by that of M.*

Proof. Denote by Γ the automorphism group of $p\colon \widetilde{M} \to M$ and by $\{\widetilde{P}_1, ..., \widetilde{P}_l\}$ the Seifert pieces of $(\widetilde{M}, \widetilde{\mathcal{T}})$. For each $i = 1, ..., l$ we know that $\widetilde{P}_i$ is homeomorphic to a product $\widetilde{F}_i \times \mathbf{S}^1$. Let $\widetilde{\alpha}$ be an element of $\widehat{H}_2^{l_1,+}(\widetilde{M}^{\widetilde{\mathcal{T}}}; \mathbf{R})$ and denote by $\mathrm{Av}(\widetilde{\alpha})$ the class obtained by *averaging* $\widetilde{\alpha}$ defined by

$$\mathrm{Av}(\widetilde{\alpha}) = \sum_{g \in \Gamma} g_\sharp(\widetilde{\alpha}) \in \Gamma \widehat{H}_2^{l_1}(\widetilde{M}; \mathbf{R})$$

By definition we have $\widetilde{\alpha} = \sum_{i \in I} \xi_i \widehat{\Theta}_{\widetilde{P}_i}(\alpha_{\widetilde{P}_i})$ where I is a subset of $\{1, ..., l\}$ and $\xi_i \in \mathbf{R}_+$ and thus since each $g \in \Gamma$ preserves the torus splitting by the naturality property of Lemma 2.1 we get

$$\mathrm{Av}(\widetilde{\alpha}) = \sum_{g \in \Gamma, i \in I} \xi_i \widehat{\Theta}_{g(\widetilde{P}_i)} \left(g_\sharp \left(\alpha_{\widetilde{P}_i} \right) \right)$$

Moreover notice that

$$\left\| \widehat{\Theta}_{g(\widetilde{P}_i)} g_\sharp \alpha_{\widetilde{P}_i} \right\|_1 \leq \left\| \widetilde{F}_i \right\|$$

For each $i = 1, ..., l$ denote by $\Omega_{\widetilde{P}_i}$ the bounded 2-cocycle of $\widetilde{M}$ constructed in Proposition 4.1 and denote by Ω the sum $\sum_{i=1}^{l} \Omega_{\widetilde{P}_i}$. Since each $g \in \Gamma$ acts one $\widetilde{M}$ as an orientation preserving diffeomorphism which preserves the orientation of the fibers (by the framing hypothesis) and the torus decomposition $\widetilde{\mathcal{T}}$ of $\widetilde{M}$ then we get

$$\langle [\Omega], \mathrm{Av}(\widetilde{\alpha}) \rangle = \mathrm{Card}(\Gamma) \sum_{i \in I} \xi_i \mathrm{Area}(\widetilde{F}_i) \leq \pi \|\mathrm{Av}(\widetilde{\alpha})\|_1$$

This proves that

$$\|\mathrm{Av}(\widetilde{\alpha})\|_1 = \mathrm{Card}(\Gamma) \sum_{i \in I} \xi_i \left\| \widetilde{F}_i \right\|$$

Recall that since $\mathrm{Av}(\widetilde{\alpha}) \in \Gamma H_2^{l_1}(\widetilde{M}; \mathbf{R})$ then by Lemma 3.2 $\|p_\sharp(\mathrm{Av}(\widetilde{\alpha}))\|_1 = \|\mathrm{Av}(\widetilde{\alpha})\|_1$. On the other hand we have

$$\|p_\sharp(\mathrm{Av}(\widetilde{\alpha}))\|_1 \leq \sum_{g \in \Gamma} \|p_\sharp g_\sharp(\widetilde{\alpha})\|_1 \leq \sum_{g \in \Gamma} \|g_\sharp(\widetilde{\alpha})\|_1 \leq \mathrm{Card}(\Gamma) \sum_{i \in I} \xi_i \left\| \widetilde{F}_i \right\|$$

We deduce that $\sum_{g\in\Gamma}\left\|p_\sharp g_\sharp(\widetilde{\alpha})\right\|_1 = \sum_{g\in\Gamma}\left\|g_\sharp(\widetilde{\alpha})\right\|_1$. On the other hand, since we know that $\left\|p_\sharp g_\sharp(\widetilde{\alpha})\right\|_1 \leq \left\|g_\sharp(\widetilde{\alpha})\right\|_1$ for any $g\in\Gamma$ then we get in particular $\left\|p_\sharp(\widetilde{\alpha})\right\|_1 = \left\|\widetilde{\alpha}\right\|_1$. □

Proof of Theorem 2.3. Let $p\colon \widetilde{M} \to M$ be a finite regular covering of M endowed with a torus splitting defined by $\widetilde{\mathcal{T}} = p^{-1}(\mathcal{T})$ and such that each component of $\widetilde{M}\setminus\widetilde{\mathcal{T}}$ is a product (such a covering exists by [LW, Proposition 4.4]). For each Seifert piece P of $M\setminus\mathcal{T}$ denote by $\widetilde{P}$ a component over P in $\widetilde{M}$. Let $\alpha\in\widehat{H}_2(P,\partial P;\mathbf{R})$. Then there exists $\widetilde{\alpha}\in\widehat{H}_2(\widetilde{P},\partial\widetilde{P};\mathbf{R})$ such that $p_\sharp(\widetilde{\alpha}) = \alpha$. Hence we have, using Lemmas 5.1 and 5.2, $\|\widehat{\Theta}_P(\alpha)\|_1 = \|\widehat{\Theta}_P(p_\sharp(\widetilde{\alpha}))\|_1 = \|p_\sharp\widehat{\Theta}_{\widetilde{P}}(\widetilde{\alpha})\|_1 = \|\widehat{\Theta}_{\widetilde{P}}(\widetilde{\alpha})\|_1 = \|\widetilde{\alpha}\|_1 \geq \|\alpha\|_1$. Indeed replacing $\widetilde{\alpha}$ by $-\widetilde{\alpha}$ we may assume that $\widehat{\Theta}_{\widetilde{P}}(\widetilde{\alpha})\in\widehat{H}_2^{l_1,+}(\widetilde{M}^{\widetilde{\mathcal{T}}})$. Since $\widehat{\Theta}_P$ is a contraction by Lemma 2.1 this proves the isometry. To complete the proof of point (i) it is sufficient to check that if α is the class of a connected horizontal surface $\mathcal{F}$ in P then $\|\widehat{\Theta}_P(\alpha)\|_1 = \|\mathcal{F}\|$. Denote by $\widetilde{\mathcal{F}}$ a component over $\mathcal{F}$ in $\widetilde{P}\subset\widetilde{M}$ and denote by n the degree of the induced covering map $\widetilde{\mathcal{F}}\to\mathcal{F}$. Let $\widetilde{\alpha}\in\widehat{H}_2(\widetilde{P},\partial\widetilde{P})$ be the integral class corresponding to $\widetilde{\mathcal{F}}$. Then we know from Lemma 5.1 that $\|\widehat{\Theta}_{\widetilde{P}}(\widetilde{\alpha})\|_1 = \|\widetilde{\mathcal{F}}\| = \|\widetilde{\alpha}\|_1$. On the other hand we know that $p_\sharp(\widehat{\Theta}_{\widetilde{P}}(\widetilde{\alpha})) = n\widehat{\Theta}_P(\alpha)$. Then we get using Lemma 5.2

$$\left\|\widetilde{\mathcal{F}}\right\| = \left\|\widetilde{\alpha}\right\|_1 = \left\|\widehat{\Theta}_{\widetilde{P}}(\widetilde{\alpha})\right\|_1 = \left\|p_\sharp(\widehat{\Theta}_{\widetilde{P}}(\widetilde{\alpha}))\right\|_1 = |n|\left\|\widehat{\Theta}_P(\alpha)\right\|_1 \leq |n|\|\alpha\|_1$$

Since $\|\alpha\|_1 \leq \|\mathcal{F}\|$ and $\|\widetilde{\mathcal{F}}\| = n\|\mathcal{F}\|$ then we get $\|\widehat{\Theta}_P(\alpha)\|_1 = \|\alpha\|_1 = \|\mathcal{F}\|$. This completes the proof of point (i).

We check the additivity property of the l_1-norm. For each P_i denote by F_i a minimal surface and for each $i = 1,...,l$ choose an element $\alpha_i\in\widehat{H}_2^+(P_i,\partial P_i;\mathbf{R})$ and we denote by α the l_1-class given by $\sum_{i=1}^l\widehat{\Theta}_{P_i}(\alpha_i)$. Denote by $\xi_1,...,\xi_l$ the non-negative real numbers such that for each $i = 1,...,l$ we have $\alpha_i = \xi_i\alpha_{P_i}$, where α_{P_i} denotes the class of F_i in $\widehat{H}_2(P_i,\partial P_i)$. For each i there exists an element $\widetilde{\alpha}_i$ in $\widehat{H}_2^+(\widetilde{P}_i,\partial\widetilde{P}_i;\mathbf{R})$ such that $p_\sharp(\widetilde{\alpha}_i) = \alpha_i$. Denote by $\widetilde{\alpha}$ the element $\sum_{i=1}^l\widehat{\Theta}_{\widetilde{P}_i}(\widetilde{\alpha}_i)\in\widehat{H}_2^{l_1,+}(\widetilde{M}^{\widetilde{\mathcal{T}}};\mathbf{R})$ such that $p_\sharp(\widetilde{\alpha}) = \alpha$. Since we may also assume that $\widehat{\Theta}_{\widetilde{P}_i}(\widetilde{\alpha}_i)\in\widehat{H}_2^{l_1,+}(\widetilde{M}^{\widetilde{\mathcal{T}}})$ for each $i = 1,...,l$ it follows from Lemma 5.2 combined with Lemma 5.1 that

$$\|\alpha\|_1 = \|\widetilde{\alpha}\|_1 = \sum_i\|\widehat{\Theta}_{\widetilde{P}_i}(\widetilde{\alpha}_i)\|_1 = \sum_i\|p_\sharp\widehat{\Theta}_{\widetilde{P}_i}(\widetilde{\alpha}_i)\|_1 = \sum_i\|\widehat{\Theta}_{P_i}(\alpha_i)\|_1$$

This completes the proof of Theorem 2.3. □

Proof of Proposition 2.4. Let $p\colon \widetilde{M}\to M$ denote a finite covering and let $q\colon\widehat{M}\to\widetilde{M}$ be a finite covering endowed with a torus splitting defined by $\widehat{\mathcal{T}} = q^{-1}(\widetilde{\mathcal{T}})$ such that each Seifert piece of $(\widehat{M},\widehat{\mathcal{T}})$ is homeomorphic to a product and such that $r = p\circ q$ is regular. Let $\{\widetilde{P}_1,...,\widetilde{P}_l\}$ denote a collection of Seifert pieces of $(\widetilde{M},\widetilde{\mathcal{T}})$ and for each i denote by $\widetilde{\alpha}_i$ an element of $\widehat{H}_2^+(\widetilde{P}_i,\partial\widetilde{P}_i;\mathbf{R})$. Let $\widetilde{\alpha}$ denote the l_1-class equal to $\sum_{i=1}^l\widehat{\Theta}_{\widetilde{P}_i}(\widetilde{\alpha}_i)$. Then using the above construction we know that there exists $\widehat{\alpha}\in\widehat{H}_2^{l_1,+}(\widehat{N}^{\widehat{\mathcal{T}}};\mathbf{R})$ such that $q_\sharp(\widehat{\alpha}) = \widetilde{\alpha}$ and $\|\widehat{\alpha}\|_1 = \|r_\sharp(\widehat{\alpha})\|_1$ by Lemma 5.2. Thus we get $\|p_\sharp\widetilde{\alpha}\|_1 = \|p_\sharp q_\sharp\widehat{\alpha}\|_1 = \|r_\sharp\widehat{\alpha}\|_1 = \|\widehat{\alpha}\|_1 \geq \|\widetilde{\alpha}\|_1$. Since any continuous map induces a contraction between l_1-homology groups we deduce that $\|p_\sharp\widetilde{\alpha}\|_1 = \|\widetilde{\alpha}\|_1$ which completes the proof of the proposition. □

6. CHARACTERIZATIONS OF FINITE COVERING MAPS

We begin this section with some recall on the Jaco Shalen Johannson torus decomposition of 3-manifolds which will be used throughout the proof of Theorem 1.1 and Corollary 1.2. Given a closed irreducible orientable 3-manifold N we denote by $\mathcal{T}_N$ the Jaco-Shalen-Johannson family of canonical tori of N and by $\mathcal{H}(N)$ (resp. $\mathcal{S}(N)$) the disjoint union of the hyperbolic (resp. Seifert) components of $N \setminus \mathcal{T}_N \times (-1,1)$ so that $N \setminus \mathcal{T}_N \times (-1,1) = \mathcal{H}(N) \cup \mathcal{S}(N)$, where $\mathcal{T}_N \times [-1,1]$ is identified with a regular neighborhood of $\mathcal{T}_N$ in such a way that $\mathcal{T}_N \simeq \mathcal{T}_N \times \{0\}$ (see [JS], [J] and [Th1]). On the other hand, we denote by $\Sigma(N) = (\Sigma(N), \emptyset)$ the *characteristic Seifert pair of* N in the sense of [JS] and [J]. We start by recalling a main consequence of the Characteristic Pair Theorem of W. Jaco and P. Shalen (see [JS, Chapter V]) which allows to control a nondegenerate map from a Seifert fibered space into an irreducible 3-manifold. We first give the definition of degenerate maps in the sense of W. Jaco and P. Shalen.

Definition 6.1. Let (S,F) be a connected Seifert pair, and let (N,T) be a connected 3-manifold pair. A map $f\colon (S,F) \to (N,T)$ is said to be *degenerate* if either

(0) the map f is inessential as a map of pairs (i.e. f is homotopic, as a map of pairs, to a map g such that $g(S) \subset T$), or

(1) the group $\mathrm{Im}(f_*\colon \pi_1 S \to \pi_1 N) = \{1\}$, or

(2) the group $\mathrm{Im}(f_*\colon \pi_1 S \to \pi_1 N)$ is cyclic and $F = \emptyset$, or

(3) the map $f|\gamma$ is homotopic in N to a constant map for some fiber γ of (S,F).

Then the Characteristic Pair Theorem of Jaco and Shalen implies the following result.

Theorem 6.2. *[Jaco, Shalen] If f is a nondegenerate map of a Seifert pair $(S,\emptyset)$ into a closed irreducible orientable 3-manifold $(N,\emptyset)$, then there exists a map f_1 of S into N, homotopic to f, such that $f_1(S) \subset \mathrm{int}(\Sigma(N))$.*

In order to prove Theorem 1.1 we first check the following technical result.

Proposition 6.3. *Let M be a closed aspherical oriented 3-manifold. Any π_1-surjective nonzero degree map $f\colon M \to N$ into a closed irreducible orientable 3-manifold satisfying the following conditions*

(i) Each Seifert component of $M \setminus \mathcal{T}_M$, resp. of $N \setminus \mathcal{T}_N$, is homeomorphic to a product, resp. to a $\mathbf{S}^1$-bundle over an orientable surface, each Seifert component of $M \setminus \mathcal{T}_M$ has at least two boundary components (if $\mathcal{T}_M \neq \emptyset$) and each component of $\mathcal{T}_M$ is shared by two distinct components of $M \setminus \mathcal{T}_M$,

(ii) $\|f_\sharp[M]\|_1 = \|[M]\|_1$, where $[M] \in H_3(M;\mathbf{R})$ is the fundamental class

(iii) $\|f_\sharp\widehat{\Theta}_P(\alpha)\|_1 = \|\widehat{\Theta}_P\alpha\|_1$ for each $\alpha \in \widehat{H}_2(P,\partial P)$ when P runs over the Seifert components of $M \setminus \mathcal{T}_M$

is homotopic to a homeomorphism.

6.1. **Proof of Proposition 6.3.** Throughout this section we always assume that the map $f\colon M \to N$ and the manifolds M, N satisfy the hypothesis of Proposition 6.3. Notice that we may assume in addition that M is not a virtual torus bundle by [W]. Thus since each Seifert piece P of M is homeomorphic to a product we may assume that P is a $\mathbf{H}^2 \times \mathbf{R}$-manifold. Hence this implies, using hypothesis (ii) and (iii), that either $\|N\| \neq 0$ or $\widehat{H}_2^{l_1}(N;\mathbf{R}) \neq \{0\}$. Hence either N has a non-empty JSJ-splitting or N admits a geometry $\mathbf{H}^3, \mathbf{H}^2 \times \mathbf{R}$ or $\widetilde{\mathrm{SL}}(2,\mathbf{R})$. The proof of Proposition 6.3 will come from the following sequence of claims.

Claim 6.4. *Let P be an incompressible $\mathbf{H}^2 \times \mathbf{R}$-submanifold of M and let c be a simple closed curve in some component T of ∂P. If there exists a horizontal surface in P whose c is a boundary component then $f_*([c]) \neq 1$ in $\pi_1 N$. Moreover, $f_*(\pi_1 P)$ is a non-abelian group.*

Proof. Denote by F a horizontal surface in P whose c is a boundary component. Since P is a $\mathbf{H}^2 \times \mathbf{R}$-manifold then F is necessarily a hyperbolic surface. In particular F has a positive simplicial volume. Suppose that $f_*([c]) = 1$. Denote by $T \times [-1, 1]$ a regular neighborhood of T such that $T = T \times \{0\}$ and parametrize $T = \mathbf{S}^1 \times \mathbf{S}^1$ such that $c = \mathbf{S}^1 \times \{*\}$. As in [Ro], consider the relation $\sim$ on M defined by $z \sim z'$ iff $z = z'$ or $z = (x, y, t) \in T \times I$, $z' = (x', y', t') \in T \times I$ and $y = y'$, $t = t'$. Denote by $X = M/\sim$ the quotient space and by $\pi\colon\ M \to X$ the quotient map. Then the map f factors throught X. Denote by $g\colon\ X \to N$ the map such that $f = g \circ \pi$. Denote by $\widehat{P}$ the image of P under π. Topologically $\widehat{P}$ is obtained from P after Dehn filling along T, identifying the meridian of a solid torus V to c so that the Seifert fibration of P extends to a Seifert fibration of $\widehat{P}$ and the image $\widehat{F}$ of F is a surface in $\widehat{P}$ obtained from F after gluing a 2-disk along each component of ∂F parallel to c. Denote by $\mathcal{C}$ the union of the components of ∂F parallel to c. Note that it follows from our construction that

$$\pi_\sharp(\widehat{\Theta}_P([F])) = \widehat{\Theta}_{\widehat{P}}([\widehat{F}]) \in H_2^{l_1}(\widehat{P}; \mathbf{R})$$

Indeed denote by z_F a relative cycle in F representing the fundamental class of F and denote by u_F a l_1-chain in ∂F such that $\partial u_F = -\partial z_F$. Then have the following equalities: $\pi_\sharp(\widehat{\Theta}_P([F])) = \pi_\sharp([z_F + u_F]) = [\pi_\sharp(z_F) + \pi_\sharp(u_F)] \in \widehat{H}_2^{l_1}(\widehat{P})$. Next consider the induces homomorphism

$$\pi_\sharp\colon\ H_2(F, \partial F) \to H_2(F/\mathcal{C}, \partial F/\mathcal{C}) = H_2(\widehat{F}, \partial\widehat{F} \coprod X_0)$$

where $X_0 = \pi(\mathcal{C}) = \mathcal{C}/\mathcal{C}$. It follows easily from the Excision Theorem that $\pi_\sharp$ is actually an isomorphism so that $\pi_\sharp(z_F)$ represents a generator of $H_2(\widehat{F}, \partial\widehat{F} \coprod X_0)$. On the other hand, there exists a 2-chain u_0 in X_0 such that $\pi_\sharp(z_F) + u_0$ is a relative cycle in $(\widehat{F}, \partial\widehat{F})$. Note that $[\pi_\sharp(z_F) + u_0] = [\pi_\sharp(z_F)]$ in $H_2(\widehat{F}, \partial\widehat{F} \coprod X_0)$. Since the inclusion $(\widehat{F}, \partial\widehat{F}) \hookrightarrow (\widehat{F}, \partial\widehat{F} \coprod X_0)$ induces an isomorphism $H_2(\widehat{F}, \partial\widehat{F}) \to H_2(\widehat{F}, \partial\widehat{F} \coprod X_0)$ then $\pi_\sharp(z_F) + u_0$ represents a generator of $H_2(\widehat{F}, \partial\widehat{F})$. Denote by $\widehat{u}_F$ a l_1-chain in $\partial\widehat{F}$ such that $\partial\widehat{u}_F = -\partial(\pi_\sharp(z_F) + u_0)$. Then since each component of $\pi(\partial P)$ has an amenable fundamental group then $[\pi_\sharp(z_F) + \pi_\sharp(u_F)] = [\pi_\sharp(z_F) + u_0 + \widehat{u}_F]$ in $H_2^{l_1}(\widehat{P}; \mathbf{R})$.

Since $\pi_\sharp(z_F) + u_0$ represents a generator of $H_2(\widehat{F}, \partial\widehat{F})$ then it follows that $\pi_\sharp(\widehat{\Theta}_P([F])) = \widehat{\Theta}_{\widehat{P}}([\widehat{F}]) \in H_2^{l_1}(\widehat{P}; \mathbf{R})$. Finally we deduce the following equalities:

$$\|\widehat{\Theta}_P([F])\|_1 \geq \|\pi_\sharp\widehat{\Theta}_P([F])\|_1 = \|\widehat{\Theta}_{\widehat{P}}([\widehat{F}])\|_1 \geq \|f_\sharp\widehat{\Theta}_P([F])\|_1 = \|\widehat{\Theta}_P([F])\|_1$$

Thus we get the following equalities:

$$\|\widehat{F}\| \geq \|\widehat{\Theta}_{\widehat{P}}([\widehat{F}])\|_1 = \|\widehat{\Theta}_P([F])\|_1 = \|F\|$$

A contradiction. This proves the first statement of the lemma.

It remains to check that $f_*(\pi_1 P)$ is a non-abelian group. Assume that $f_*(\pi_1 P)$ is an abelian subgroup of $\pi_1 N$. Then necessarily $f_*(\pi_1 P)$ is isomorphic to a free abelian group of rank ≤ 3. Denote by X a $K(f_*(\pi_1 P), 1)$-space (X is homeomorphic to either $\mathbf{D}^3, \mathbf{S}^1 \times \mathbf{D}^2, \mathbf{S}^1 \times \mathbf{S}^1 \times I$ or $\mathbf{S}^1 \times \mathbf{S}^1 \times \mathbf{S}^1$). Then the map $f|P\colon\ P \to N$ factors through

X so that we have the following commutative diagram

$$\begin{tikzcd}
\widehat{H}_2(P,\partial P) \arrow[r, "\widehat{\Theta}_P"] \arrow[d, "\simeq"] & \widehat{H}_2^{l_1}(P) \arrow[r, "i_\sharp"] \arrow[dr] & \widehat{H}_2^{l_1}(M) \arrow[r, "f_\sharp"] & \widehat{H}_2^{l_1}(N) \\
\mathbf{R} & & \widehat{H}_2^{l_1}(X) \arrow[ur] &
\end{tikzcd}$$

Since $\widehat{H}_2^{l_1}(X)$ is trivial (because X has an amenable fundamental group) then we get a contradiction with hypothesis (iii) of Proposition 6.3. $\square$

Claim 6.5. *Let $P = F \times S^1$ be a 3-manifold such that F is an orientable hyperbolic surface with non connected boundary. Then for any simple closed curve c of ∂P that is not homotopic to the fiber of P, there exists a horizontal surface $(H, \partial H)$ in $(P, \partial P)$ such that c is parallel to a component of ∂H.*

Proof. Denote by T_1 the component of ∂P which contains c and by $T_2, ..., T_r$ the other components of ∂P with $r \geq 2$. For each $i = 1, ..., r$ fix a basis $\langle s_i, h\rangle$, where s_i is a section of T_i with respect to the $\mathbf{S}^1$-fibration of P such that $s_1 + ... + s_r$ is nul-homologous in P and where h denotes the fiber of P. Denote by (α, β) the coprime integers with $\alpha \neq 0$ such that $c = \alpha[s_1] + \beta[h]$. Then

$$[c] + \alpha[s_2] + ... + \alpha[s_r] - \beta[h] = 0 \text{ in } H_1(P; \mathbf{Z})$$

Thus there exists a nontrivial class η in $H_2(P, \partial P; \mathbf{Z})$ such that

$$\partial\eta = ((\alpha, \beta), (\alpha, 0), ..., (\alpha, 0), (\alpha, -\beta))$$

in $H_1(\partial P) = H_1(T_1) \oplus H_1(T_2) \oplus ... \oplus H_1(T_{r-1}) \oplus H_1(T_r)$. Denote by H an incompressible surface representing η. Then H is necessarily a horizontal surface and thus c is parallel to some components of ∂H. This proves the claim. $\square$

Claim 6.6. *The map $f|T\colon\ T \to N$ is π_1-injective for any characteristic torus T in M.*

Proof. Let T be a characteristic torus of M. From the Rigidity Theorem of Soma [So] and from hypothesis (i) it is sufficient to consider the case where T is shared by two distinct Seifert components Σ_1 and Σ_2 of $M \setminus \mathcal{T}_M$. For each $i = 1, 2$, denote by h_i the $\mathbf{S}^1$-fiber of Σ_i. Combining Claims 6.4 and 6.5 we deduce that if $f|T\colon\ T \to N$ is not π_1-injective then $f_*(h_i) = \{1\}$. Since h_1 and h_2 generate a rank 2 subgroup of $\pi_1 T$ (by minimality of the JSJ-decomposition) we get a contradiction. This proves the claim.

$\square$

Claim 6.7. *There is a map g homotopic to f such that for each Seifert piece of Σ of N then each component of $g^{-1}(\Sigma)$ is a Seifert piece of M.*

Proof. By Lemma 6.6 one can apply the Theorem 6.2 combined with [So, Rigidity Theorem] which imply that one can arrange f by a homotopy so that for each canonical torus U of N then $f^{-1}(U)$ is a disjoint union of canonical tori of M. Hence for each Seifert piece Σ of N the space $f^{-1}(\Sigma)$ is a canonical graph submanifold of M (i.e. a submanifold which is the union of some Seifert pieces of M).

If a component G of $f^{-1}(\Sigma)$ is not a Seifert manifold then there exists a canonical torus T in the interior of G which is shared by two distinct Seifert pieces Σ_1 and Σ_2 of G. Since by Claim 6.4 $f_*(\pi_1\Sigma_1)$ is not abelian then using [JS, Addendum to Theorem VI.I.6] we know that $f|\Sigma_1$ is homotopic to a fiber preserving map. Since $f|T$ is π_1-injective we get a contradiction by the minimality of the JSJ-decomposiiton. This proves the claim. $\square$

By hypothesis (ii) one can apply [So, Rigidity Theorem]. Thus one may assume that f induces a $\deg(f)$-covering map from $\mathcal{H}(M)$ to $\mathcal{H}(N)$. Moreover since f is π_1-surjective then to complete the proof of Proposition 6.3 it remains to check the following

Claim 6.8. *There is a map g homotopic to f, rel. to $\mathcal{H}(M)$, such that for each Seifert piece Σ of N and for each component G of $g^{-1}(\Sigma)$ then $g|G\colon\ G \to \Sigma$ is a covering map.*

Proof. First of all note that for each component G of $f^{-1}(\Sigma)$ then $f|G\colon\ G \to \Sigma$ is non-degenerate and fiber preserving.

Indeed if ∂G is non-empty this comes from Claim 6.6. If $\partial G = \emptyset$ then $M = G$ and $N = \Sigma$ so that $f\colon\ M \to N$ is a nonzero degree map between $\mathbf{S}^1$-bundles over orientable hyperbolic surfaces. Since $\deg(f) \neq 0$ then $f_*(h) \neq 1$, where h denotes the fiber of M and by Claim 6.4, $f_*(\pi_1 M)$ is a non-abelian group.

On the other hand, notice that Σ is necessarily homeomorphic to a product.

Indeed if $\partial\Sigma \neq \emptyset$ this is obvious and if $\partial\Sigma = \emptyset$ this comes from the following argument: if Σ is not homeomorphic to a product then the bundle has a non-zero Euler number and using the Seifert volume in [Br-G1, Theorem 3 and Lemma 3] and in [Br-G2, Theorem 4] we get a contradiction (since G has a zero Euler number and $\deg(f) \neq 0$).

On the other hand notice that for each component G of $f^{-1}(\Sigma)$ then $\deg(f|G\colon\ G \to \Sigma) \neq 0$. Indeed, suppose that $\deg(f|G\colon\ G \to \Sigma) = 0$. Since by construction $f|G$ is an allowable map (in the sense of [Ro]) then it induces a zero degree map $\pi\colon\ K \to F$ where K, resp. F, is a hyperbolic surface such that $G = K \times \mathbf{S}^1$, resp. $\Sigma = F \times \mathbf{S}^1$. Let $\mathcal{F}$ denote a component of $(f|G)^{-1}(F)$. Arrange f so that $\mathcal{F}$ is incompressible in G. Since f is non-degenerate and fiber preserving then $\mathcal{F}$ is necessarily a horizontal surface. Since $f|\mathcal{F}\colon\ \mathcal{F} \to F$ factors throught π then $\deg(f|\mathcal{F}\colon\ \mathcal{F} \to F) = 0$. Thus using the naturality property we get the equality

$$f_\sharp(\widehat{\Theta}_G[\mathcal{F}]) = \widehat{\Theta}_\Sigma f_\sharp[\mathcal{F}] = 0$$

This gives a contradiction with hypothesis (iii) of Proposition 6.3 since

$$\|\widehat{\Theta}_G[\mathcal{F}]\|_1 = \|\mathcal{F}\| > 0$$

Hence since $\deg(f|G\colon\ G \to \Sigma) \neq 0$ then f induces a nonzero degree map $f|\mathcal{F}\colon\ \mathcal{F} \to F$ so that we get

$$f_\sharp(\widehat{\Theta}_G[\mathcal{F}]) = \widehat{\Theta}_\Sigma f_\sharp[\mathcal{F}] = \deg(f|\mathcal{F}\colon\ \mathcal{F} \to F)\widehat{\Theta}_\Sigma[F]$$

which implies that

$$\|\mathcal{F}\| = |\deg(f|\mathcal{F}\colon\ \mathcal{F} \to F)| \times \|F\|$$

Thus we get the equality

$$\|K\| = \deg(\pi) \times \|F\|$$

Hence π is homotopic to a covering map which implies that $f|G$ is also homotopic to a covering map. This proves the claim and completes the proof of the Proposition 6.3. □

6.2. **Proof of Theorem 1.1.** First of all note that according to [W], [So] and the hypothesis of the theorem, we may assume that either $\mathcal{T}_M \neq \emptyset$ or M is a $\mathbf{H}^2 \times \mathbf{R}$-manifold. Thus we have $\widehat{H}_2^{l_1}(M;\mathbf{R}) \neq 0$. Then either N has a non-empty JSJ-decomposition or N admits one of the following geometry: $\mathbf{H}^2 \times \mathbf{R}$ or $\widetilde{\mathrm{SL}}(2,\mathbf{R})$.

Denote by $P_1, ..., P_k$ the Seifert pieces of M. Then $[M]^{\mathcal{T}_M} = \widehat{\Theta}_{P_1}\alpha_{P_1} + ... + \widehat{\Theta}_{P_k}\alpha_{P_k}$. Then using the additivity property of the l_1-norm and the isometry hypothesis we have

$$\sum_{i=1}^{k}\left\|\widehat{\Theta}_{P_i}\alpha_{P_i}\right\|_1 \geq \sum_{i=1}^{k}\left\|f_\sharp\widehat{\Theta}_{P_i}\alpha_{P_i}\right\|_1 \geq \left\|f_\sharp\sum_{i=1}^{k}\widehat{\Theta}_{P_i}\alpha_{P_i}\right\|_1 = \left\|\sum_{i=1}^{k}\widehat{\Theta}_{P_i}\alpha_{P_i}\right\|_1$$

and since, by the additivity property of Theorem 2.3,

$$\left\|\sum_{i=1}^{k}\widehat{\Theta}_{P_i}\alpha_{P_i}\right\|_1 = \sum_{i=1}^{k}\left\|\widehat{\Theta}_{P_i}\alpha_{P_i}\right\|_1$$

hence we get

$$\sum_{i=1}^{k}\left\|f_\sharp\widehat{\Theta}_{P_i}\alpha_{P_i}\right\|_1 = \sum_{i=1}^{k}\left\|\widehat{\Theta}_{P_i}\alpha_{P_i}\right\|_1$$

and thus $\|f_\sharp\widehat{\Theta}_{P_i}\alpha_{P_i}\|_1 = \|\widehat{\Theta}_{P_i}\alpha_{P_i}\|_1$ for any $i = 1, ..., k$ which implies, using Lemma 2.2, that $\|f_\sharp\widehat{\Theta}_P\alpha\|_1 = \|\widehat{\Theta}_P\alpha\|_1$ for each $\alpha \in \widehat{H}_2(P, \partial P)$ and P in $\mathcal{S}(M)$. Consider now the following commutative diagram

$$\begin{tikzcd}
M_2 \arrow[r, "f_2"] \arrow[d, "q"'] & N_2 \arrow[d, "p"] \\
M_1 \arrow[r, "f_1"] \arrow[d, "s"'] & N_1 \arrow[d, "r"] \\
M \arrow[r, "f"] & N
\end{tikzcd}$$

obtained as follows. The map $s\colon M_1 \to M$ is a finite covering such that each Seifert piece of M_1 is a product with at least two boundary components, if $\mathcal{T}_M \neq \emptyset$, and each canonical torus of M_1 is shared by two distinct components of $M_1 \setminus \mathcal{T}_{M_1}$ (for the existence of such a covering see [DW, Lemmas 3.2 and 3.5]), the map $r\colon N_1 \to N$ is a finite covering corresponding to the subgroup $f_*s_*(\pi_1 M_1)$ in $\pi_1 N$, which is of finite index since $\deg(f) \neq 0$, the map $f_1\colon M_1 \to N_1$ is a lifting of $f \circ s$ which exists by our construction, the map $p\colon N_2 \to N_1$ is a finite covering such that each Seifert piece of N_2 is a $\mathbf{S}^1$-bundle over an orientable surface and $f_2\colon M_2 \to N_2$ is the finite covering of f_1 corresponding to p, and $q\colon M_2 \to M_1$ is the covering corresponding to the subgroup $(f_1)_*^{-1}(p_*\pi_1 N_2)$. Notice that it follows from the construction that f_1 and f_2 are π_1-surjective. On the other and let α be an element of $\widehat{H}_2(P, \partial P)$, where P is a Seifert piece of M_2. Then using the isometric properties of finite coverings of Proposition 2.4 together with the commutativity of the diagram we get the following equalities (**)

$$\left\|\widehat{\Theta}_P\alpha\right\|_1 = \left\|f_\sharp s_\sharp q_\sharp\widehat{\Theta}_P\alpha\right\|_1 = \left\|r_\sharp p_\sharp (f_2)_\sharp\widehat{\Theta}_P\alpha\right\|_1 \leq \left\|(f_2)_\sharp\widehat{\Theta}_P\alpha\right\|_1 \leq \left\|\widehat{\Theta}_P\alpha\right\|_1$$

Indeed replacing α by $-\alpha$ we may assume that $\widehat{\Theta}_P\alpha \in \widehat{H}_2^{l_1,+}\left(M_2^{\mathcal{T}_{M_2}}\right)$ and since the covering maps q and s preserve the torus decompositions then $q_\sharp\widehat{\Theta}_P\alpha \in \widehat{H}_2^{l_1,+}\left(M_1^{\mathcal{T}_{M_1}}\right)$ and $s_\sharp q_\sharp\widehat{\Theta}_P\alpha \in \mathrm{Im}(\widehat{\Theta}_Q)$ for some Seifert piece Q of M. Thus one can apply Proposition 6.3 which implies that f_2 is homotopic to a homeomorphism. Since p, q, r, s are finite covering maps then this implies that f is π_1-injective. Consider the finite covering $\widetilde{N} \to N$ corresponding to $f_*(\pi_1 M)$. Then f lifts to a map $\widetilde{f}\colon M \to \widetilde{N}$ inducing an isomorphism at the π_1-level. We deduce from this point using [Wa] and [O, Section 5.3, Theorem 6] that $\widetilde{f}$ is a homeomorphism. This implies that f is a covering map and completes the proof of Theorem 1.1.

6.3. **Proof of Corollary 1.2.** Using the duality we may assume that f induces an isometry $f_\sharp\colon\ \widehat{H}_2^{l_1}(M;\mathbf{R}) \to \widehat{H}_2^{l_1}(N;\mathbf{R})$. Then according to Theorem 1.1 it is sufficient to consider the case where M is a $\widetilde{\mathrm{SL}}(2,\mathbf{R})$-manifold. Then in particular we have $\widehat{H}_2^{l_1}(M;\mathbf{R}) \neq 0$ and thus $\widehat{H}_2^{l_1}(N;\mathbf{R}) \neq 0$. On the other hand, since $f\colon\ M \to N$ is a degree one map then N admits necessarily a geometry $\mathbf{H}^2 \times \mathbf{R}$ or $\widetilde{\mathrm{SL}}(2,\mathbf{R})$. Since f is a degree one map, then it is π_1-surjective and thus it is homotopic to a fiber preserving map and $f_*([\gamma]) \neq \{1\}$ in $\pi_1 N$ for any fiber γ of M. If N is a $\mathbf{H}^2 \times \mathbf{R}$-manifold, choose a horizontal surface H is N. Then one can arrange f by a homotopy so that $f^{-1}(H)$ is either a horizontal or a vertical surface if M. Since M is a $\widetilde{\mathrm{SL}}(2,\mathbf{R})$-manifold then there are no horizontal surface in M so that $f^{-1}(H)$ consists necessarily of vertical surfaces. A contradiction since f is a non-degenerate fiber preserving. Thus N is also a $\widetilde{\mathrm{SL}}(2,\mathbf{R})$-manifold.

Since N is a Haken manifold then there exists a non-empty torus splitting $\mathcal{U}$ of N. On the other hand note that one can arrange f by a homotopy so that $\mathcal{T} = f^{-1}(\mathcal{U})$ is also a torus splitting of M. As in paragraph 6.2, consider the commutative diagram

$$\begin{tikzcd}
(M_2,\mathcal{T}_2) \arrow[r, "f_2"] \arrow[d, "q"'] & (N_2,\mathcal{U}_2) \arrow[d, "p"] \\
(M_1,\mathcal{T}_1) \arrow[r, "f_1"] \arrow[d, "s"'] & (N_1,\mathcal{U}_1) \arrow[d, "r"] \\
(M,\mathcal{T}) \arrow[r, "f"] & (N,\mathcal{U})
\end{tikzcd}$$

where $s\colon\ M_1 \to M$ is a finite covering such that each component of $M_1 \setminus \mathcal{T}_1$ is a product with at least two boundary components, $\mathcal{T}_1 = s^{-1}(\mathcal{T})$, and each component of $\mathcal{T}_1$ is shared by two distinct components of $M_1 \setminus \mathcal{T}_1$, the map $r\colon\ N_1 \to N$ is a finite covering corresponding to the subgroup $f_*s_*(\pi_1 M_1)$ in $\pi_1 N$ and $\mathcal{U}_1 = r^{-1}(\mathcal{U})$, the map $f_1\colon\ M_1 \to N_1$ is a lifting of $f \circ s$, the map $p\colon\ N_2 \to N_1$ is a finite covering with $\mathcal{U}_2 = p^{-1}(\mathcal{U}_1)$ such that each component of $N_2 \setminus \mathcal{U}_2$ is a product and $f_2\colon\ M_2 \to N_2$ is the finite covering of f_1 corresponding to p with $\mathcal{T}_2 = q^{-1}(\mathcal{T}_1)$. Note that, using the same argument as in (**) paragraph 6.2, we obtain $\|(f_2)\sharp\widehat{\Theta}_P(\alpha)\|_1 = \|\widehat{\Theta}_P\alpha\|_1$ for each $\alpha \in \widehat{H}_2(P,\partial P)$ when P runs over the components of $M_2 \setminus \mathcal{T}_2$. Hence, since using Claim 6.4 and 6.5 the map $f_2|\mathcal{T}_2$ is π_1-injective, one can apply the proof of Claim 6.8 which implies that for each component Q of $N_2 \setminus \mathcal{U}_2$ then $f_2|f_2^{-1}(Q)\colon\ f_2^{-1}(Q) \to Q$ is a covering map. This completes the proof of Corollary 1.2.

REFERENCES

[BG] J. BARGES, E. GHYS, *Surfaces et cohomologie bornée*, Invent. Math. 92 (1988), no. 3, 509–526.

[BW] M. BOILEAU, S. WANG, *Non-zero degree maps and surface bundles over* $\mathbf{S}^1$, J. Differential Geom. 43 (1996), no. 4, 789–806.

[Br-G1] R. BROOKS, W. GOLDMAN, *Volumes in Seifert space*, Duke Math. J. 51 (1984), no. 3, 529–545.

[Br-G2] R. BROOKS, W. GOLDMAN, *The Godbillon-Vey invariant of a transversely homogeneous foliation*, Trans. Amer. Math. Soc. 286 (1984), no. 2, 651–664.

[DW] P. DERBEZ, S. WANG, *Finiteness of mapping degrees and* $\mathrm{PSL}(2,\mathbf{R})$*-volume on graph manifolds*, Preprint (2008).

[FS] K. FUJIWARA, T. SOMA, *Bounded classes in the cohomology of manifolds*, Dedicated to John Stallings on the occasion of his 65th birthday. Geom. Dedicata 92 (2002), 73–85.

[G] M. GROMOV, *Volume and bounded cohomology*, Inst. Hautes Études Sci. Publ. Math. No. 56, (1982), 5–99.

[JS] W. JACO, P.B. SHALEN, *Seifert fibered space in 3-manifolds*, Mem. Amer. Math. Soc. 21 (1979).

[J] K. JOHANNSON, *Homotopy equivalences of 3-manifolds with boundaries*, Lecture Notes in Mathematics, 761. Springer, Berlin, 1979.

[LW] J. LUECKE, Y. WU, *Relative Euler number and finite covers of graph manifolds* Geometric topology (Athens, GA, 1993), 80–103, AMS/IP Stud. Adv. Math., 2.1, Amer. Math. Soc., Providence, RI, 1997.

[MM] S. MATSUMOTO, S MORITA, *Bounded cohomology of certain groups of homeomorphisms*, Proc. Amer. Math. Soc. 94 (1985), no. 3, 539–544.

[O] P. ORLIK, *Seifert manifolds*, Lecture Notes in Mathematics, Vol. 291.

[Ro] Y. RONG, *Maps between Seifert fibered spaces of infinite* π_1, Pacific J. Math. 160 (1993), no. 1, 143–154.

[Ro1] Y. RONG, *Degree one maps of Seifert manifolds and a note on Seifert volume*, Topology Appl. 64 (1995), no. 2, 191–200.

[So] T. SOMA, *A rigidity theorem for Haken manifolds*, Math. Proc. Cambridge Philos. Soc. 118 (1995), no. 1, 141–160.

[Th] W. THURSTON, *The geometry and topology of 3-manifolds*, Lectures Notes, Princeton Univ., 1979.

[Th1] W. THURSTON, *Three dimensional manifolds, Kleinian groups and hyperbolic geometry*, Bull. Amer. Math. Soc. (N.S.) 6 (1982), no. 3, 357–381.

[Wa] F. WALDHAUSEN, *On irreducible 3-manifolds which are sufficiently large*, Ann. of Math. 87 (1968), 56-88.

[W] S. WANG, *The existence of maps of nonzero degree between aspherical 3-manifolds*, Math. Z. 208 (1991), no. 1, 147160.

[WZ] S. WANG, Q. ZHOU, *Any* 3*-manifold* 1*-dominates at most finitely many geometric* 3*-manifolds*, Math. Ann. 322 (2002), no. 3, 525-535.

LATP, UMR 6632, CENTRE DE MATHÉMATIQUES ET D'INFORMATIQUE, TECHNOPOLE DE CHATEAU-GOMBERT, 39, RUE FRÉDÉRIC JOLIOT-CURIE - 13453 MARSEILLE CEDEX 13
E-mail address: `derbez@cmi.univ-mrs.fr`